

THE TRUNCATED MOMENT PROBLEM ON REDUCIBLE CUBIC CURVES I: PARABOLIC AND CIRCULAR TYPE RELATIONS

SEONGUK YOO AND ALJAŽ ZALAR^{2,Q}

ABSTRACT. In this article we study the bivariate truncated moment problem (TMP) of degree $2k$ on reducible cubic curves. First we show that every such TMP is equivalent after applying an affine linear transformation to one of 8 canonical forms of the curve. The case of the union of three parallel lines was solved in [Zal22a], while the degree 6 cases in [Yoo17b]. Second we characterize in terms of concrete numerical conditions the existence of the solution to the TMP on two of the remaining cases concretely, i.e., a union of a line and a circle $y(ay + x^2 + y^2) = 0, a \in \mathbb{R} \setminus \{0\}$, and a union of a line and a parabola $y(x - y^2) = 0$. In both cases we also determine the number of atoms in a minimal representing measure.

1. INTRODUCTION

Given a real 2–dimensional sequence

$$\beta \equiv \beta^{(2k)} = \{\beta_{0,0}, \beta_{1,0}, \beta_{0,1}, \dots, \beta_{2k,0}, \beta_{2k-1,1}, \dots, \beta_{1,2k-1}, \beta_{0,2k}\}$$

of degree $2k$ and a closed subset K of \mathbb{R}^2 , the **truncated moment problem (K –TMP)** supported on K for $\beta^{(2k)}$ asks to characterize the existence of a positive Borel measure μ on \mathbb{R}^2 with support in K , such that

$$(1.1) \quad \beta_{i,j} = \int_K x^i y^j d\mu \quad \text{for } i, j \in \mathbb{Z}_+, 0 \leq i + j \leq 2k.$$

If such a measure exists, we say that $\beta^{(2k)}$ has a representing measure supported on K and μ is its **K –representing measure (K –rm)**.

In the degree-lexicographic order

$$1, X, Y, X^2, XY, Y^2, \dots, X^k, X^{k-1}Y, \dots, Y^k$$

of rows and columns, the corresponding moment matrix to β is equal to

$$(1.2) \quad \mathcal{M}(k) \equiv \mathcal{M}(k; \beta) := \begin{pmatrix} \mathcal{M}[0, 0](\beta) & \mathcal{M}[0, 1](\beta) & \cdots & \mathcal{M}[0, k](\beta) \\ \mathcal{M}[1, 0](\beta) & \mathcal{M}[1, 1](\beta) & \cdots & \mathcal{M}[1, k](\beta) \\ \vdots & \vdots & \ddots & \vdots \\ \mathcal{M}[k, 0](\beta) & \mathcal{M}[k, 1](\beta) & \cdots & \mathcal{M}[k, k](\beta) \end{pmatrix},$$

Date: October 27, 2024.

2020 Mathematics Subject Classification. Primary 44A60, 47A57, 47A20; Secondary 15A04, 47N40.

Key words and phrases. Truncated moment problems; K –moment problems; K –representing measure; Minimal measure; Moment matrix extensions.

²Supported by the Slovenian Research Agency program P1-0288 and grants J1-50002, J1-2453, J1-3004.

^QThis work was performed within the project COMPUTE, funded within the QuantERA II Programme that has received funding from the EU’s H2020 research and innovation programme under the GA No 101017733 .

where

$$\mathcal{M}[i, j](\beta) := \begin{pmatrix} \beta_{i+j,0} & \beta_{i+j-1,1} & \beta_{i+j-2,2} & \cdots & \beta_{i,j} \\ \beta_{i+j-1,1} & \beta_{i+j-2,2} & \beta_{i+j-3,3} & \cdots & \beta_{i-1,j+1} \\ \beta_{i+j-2,2} & \beta_{i+j-3,3} & \beta_{i+j-4,4} & \cdots & \beta_{i-2,j+2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \beta_{j,i} & \beta_{j-1,i+1} & \beta_{j-2,i+2} & \cdots & \beta_{0,i+j} \end{pmatrix}.$$

Let $\mathbb{R}[x, y]_{\leq k} := \{p \in \mathbb{R}[x, y] : \deg p \leq k\}$ stand for the set of real polynomials in variables x, y of total degree at most k . For every $p(x, y) = \sum_{i,j} a_{i,j} x^i y^j \in \mathbb{R}[x, y]_{\leq k}$ we define its **evaluation** $p(X, Y)$ on the columns of the matrix $\mathcal{M}(k)$ by replacing each capitalized monomial $X^i Y^j$ in $p(X, Y) = \sum_{i,j} a_{i,j} X^i Y^j$ by the column of $\mathcal{M}(k)$, indexed by this monomial. Then $p(X, Y)$ is a vector from the linear span of the columns of $\mathcal{M}(k)$. If this vector is the zero one, i.e., all coordinates are equal to 0, then we say p is a **column relation** of $\mathcal{M}(k)$. A column relation p is **nontrivial**, if $p \neq 0$. We denote by $\mathcal{Z}(p) := \{(x, y) \in \mathbb{R}^2 : p(x, y) = 0\}$, the zero set of p . We say that the matrix $\mathcal{M}(k)$ is **recursively generated (rg)** if for $p, q, pq \in \mathbb{R}[x, y]_{\leq k}$ such that p is a column relation of $\mathcal{M}(k)$, it follows that pq is also a column relation of $\mathcal{M}(k)$. The matrix $\mathcal{M}(k)$ is **p -pure**, if the only column relations of $\mathcal{M}(k)$ are those determined recursively by p . We call a sequence β **p -pure**, if $\mathcal{M}(k)$ is p -pure.

A **concrete solution** to the TMP is a set of necessary and sufficient conditions for the existence of a K -representing measure μ , that can be tested in numerical examples. Among necessary conditions, $\mathcal{M}(k)$ must be positive semidefinite (psd) and rg [CF04, Fia95], and by [CF96], if the support $\text{supp}(\mu)$ of μ is a subset of $\mathcal{Z}(p)$ for a polynomial $p \in \mathbb{R}[x, y]_{\leq k}$, then p is a column relation of $\mathcal{M}(k)$. The bivariate K -TMP is concretely solved in the following cases:

- (1) $K = \mathcal{Z}(p)$ for a polynomial p with $1 \leq \deg p \leq 2$.

Assume that $\deg p = 2$. By applying an affine linear transformation it suffices to consider one of the canonical cases: $x^2 + y^2 = 1$, $y = x^2$, $xy = 1$, $xy = 0$, $y^2 = y$. The case $x^2 + y^2 = 1$ is equivalent to the univariate trigonometric moment problem, solved in [CF02]. The other four cases were tackled in [CF02, CF04, CF05, Fia15] by applying the far-reaching **flat extension theorem (FET)** [CF96, Theorem 7.10] (see also [CF05b, Theorem 2.19] and [Lau05] for an alternative proof), which states that $\beta^{(2k)}$ admits a (rank $\mathcal{M}(k)$)-atomic rm if and only if $\mathcal{M}(k)$ is psd and admits a rank-preserving extension to a moment matrix $\mathcal{M}(k+1)$. For an alternative approach with shorter proofs compared to the original ones by reducing the problem to the univariate setting see [BZZ1, Section 6] (for $xy = 0$), [Zal22a] (for $y^2 = y$), [Zal22b] (for $xy = 1$) and [Zal23] (for $y = x^2$).

For $\deg p = 1$ the solution is [CF08, Proposition 3.11] and uses the FET, but can be also derived in the univariate setting (see [Zal23, Remark 3.3.(4)])

- (2) $K = \mathbb{R}^2$, $k = 2$ and $\mathcal{M}(2)$ is invertible. This case was first solved nonconstructively using convex geometry techniques in [FN10] and later on constructively in [CY16] by a novel rank reduction technique.
- (3) K is one of $\mathcal{Z}(y - x^3)$ [Fia11, Zal21], $\mathcal{Z}(y^2 - x^3)$ [Zal21], $\mathcal{Z}(y(y - a)(y - b))$ [Yoo17a, Zal22a], $a, b \in \mathbb{R} \setminus \{0\}$, $a \neq b$, or $\mathcal{Z}(xy^2 - 1)$ [Zal22b]. The main technique in [Fia11] is the FET, while in [Zal21, Zal22a, Zal22b] the reduction to the univariate TMP is applied.
- (4) $\mathcal{M}(k)$ has a special feature called *recursive determinateness* [CF13] or *extremality* [CFM08].
- (5) $\mathcal{M}(3)$ satisfies symmetric cubic column relations which can only cause extremal moment problems. In order to satisfy the variety condition, another symmetric column relation must exist, and the solution was obtained by checking consistency [CY14].

- (6) Non-extremal sextic TMPs with rank $\mathcal{M}(3) \leq 8$ and with finite or infinite algebraic varieties [CY15].
- (7) $\mathcal{M}(3)$ with reducible cubic column relations [Yoo17b].

The solutions to the K -TMP, which are not concrete in the sense of definition from the previous paragraph, are known in the cases $K = \mathcal{Z}(y - q(x))$ and $K = \mathcal{Z}(yq(x) - 1)$, where $q \in \mathbb{R}[x]$. [Fia11, Section 6] gives a solution in terms of the bound on the degree m for which the existence of a positive extension $\mathcal{M}(k + m)$ of $\mathcal{M}(k)$ is equivalent to the existence of a rm. In [Zal23] the bound on m is improved to $m = \deg q - 1$ for curves of the form $y = q(x)$, $\deg q \geq 3$, and to $m = \ell + 1$ for curves of the form $yx^\ell = 1$, $\ell \in \mathbb{N} \setminus \{1\}$.

References to some classical work on the TMP are monographs [Akh65, AK62, KN77], while for a recent development in the area we refer a reader to [Sch17]. Special cases of the TMP have also been considered in [Kim14, Ble15, Fia17, DS18, BF20, Kim21], while [Nie14] considers subspaces of the polynomial algebra and [CGIK+] the TMP for commutative \mathbb{R} -algebras.

The motivation for this paper was to solve the TMP concretely on some reducible cubic curves, other than the case of three parallel lines solved in [Zal22a]. Applying an affine linear transformation we show that every such TMP is equivalent to the TMP on one of 8 canonical cases of reducible cubics of the form $yc(x, y) = 0$, where $c \in \mathbb{R}[x, y]$, $\deg c = 2$. In this article we solve the TMP for the cases $c(x, y) = ay + x^2 + y^2$, $a \in \mathbb{R} \setminus \{0\}$, and $c(x, y) = x - y^2$, which we call the *circular* and the *parabolic type*, respectively. The main idea is to characterize the existence of a decomposition of β into the sum $\beta^{(\ell)} + \beta^{(c)}$, where $\beta^{(\ell)} = \{\beta_{i,j}^{(\ell)}\}_{i,j \in \mathbb{Z}_+, 0 \leq i+j \leq 2k}$ and $\beta^{(c)} = \{\beta_{i,j}^{(c)}\}_{i,j \in \mathbb{Z}_+, 0 \leq i+j \leq 2k}$ admit a \mathbb{R} -rm and a $\mathcal{Z}(c)$ -rm, respectively. Due to the form of the cubic $yc(x, y) = 0$, it turns out that all but two moments of $\beta^{(\ell)}$ and $\beta^{(c)}$ are not already fixed by the original sequence, i.e., $\beta_{0,0}^{(\ell)}$, $\beta_{1,0}^{(\ell)}$, $\beta_{0,0}^{(c)}$, $\beta_{1,0}^{(c)}$ in the circular type case and $\beta_{0,0}^{(\ell)}$, $\beta_{2k,0}^{(\ell)}$, $\beta_{0,0}^{(c)}$, $\beta_{2k,0}^{(c)}$ in the parabolic type case. Then, by an involved analysis, the characterization of the existence of a decomposition $\beta = \beta^{(\ell)} + \beta^{(c)}$ can be done in both cases. We also characterize the number of atoms in a minimal representing measure, i.e., a measure with the minimal number of atoms in the support.

1.1. Readers Guide. The paper is organized as follows. In Section 2 we present some preliminary results needed to establish the main results of the paper. In Section 3 we show that to solve the TMP on every reducible cubic curve it is enough to consider 8 canonical type relations (see Proposition 3.1). In Section 4 we present the general procedure for solving the TMP on all but one of the canonical types and prove some results that apply to them. Then in Sections 5 and 6 we specialize to the circular and the parabolic type relations and solve them concretely (see Theorems 5.1 and 6.1). In both cases we show, by numerical examples, that there are pure sequences $\beta^{(6)}$ with a psd $\mathcal{M}(3)$ but without a rm (see Examples 5.3 and 6.3).

2. PRELIMINARIES

We write $\mathbb{R}^{n \times m}$ for the set of $n \times m$ real matrices. For a matrix M we call the linear span of its columns a **column space** and denote it by $\mathcal{C}(M)$. The set of real symmetric matrices of size n will be denoted by S_n . For a matrix $A \in S_n$ the notation $A \succ 0$ (resp. $A \succeq 0$) means A is positive definite (pd) (resp. positive semidefinite (psd)). We write $\mathbf{0}_{t_1, t_2}$ for a $t_1 \times t_2$ matrix with only zero entries and $\mathbf{0}_t = \mathbf{0}_{t,t}$ for short, where $t_1, t_2, t \in \mathbb{N}$. The notation $E_{i,j}^{(\ell)}$, $\ell \in \mathbb{N}$, stands for the usual $\ell \times \ell$ coordinate matrix with the only nonzero entry at the position (i, j) , which is equal to 1.

In the rest of this section let $k \in \mathbb{N}$ and $\beta \equiv \beta^{(2k)} = \{\beta_{i,j}\}_{i,j \in \mathbb{Z}_+, 0 \leq i+j \leq 2k}$ be a bivariate sequence of degree $2k$.

2.1. Moment matrix. Let $\mathcal{M}(k)$ be the moment matrix of β (see (1.2)). Let Q_1, Q_2 be subsets of the set $\{X^i Y^j : i, j \in \mathbb{Z}_+, 0 \leq i+j \leq k\}$. We denote by $(\mathcal{M}(k))_{Q_1, Q_2}$ the submatrix of $\mathcal{M}(k)$ consisting of the rows indexed by the elements of Q_1 and the columns indexed by the elements of Q_2 . In case $Q := Q_1 = Q_2$, we write $(\mathcal{M}(k))_Q := (\mathcal{M}(k))_{Q, Q}$ for short.

2.2. Affine linear transformations. The existence of representing measures is invariant under invertible affine linear transformations of the form

$$(2.1) \quad \phi(x, y) = (\phi_1(x, y), \phi_2(x, y)) := (a + bx + cy, d + ex + fy), \quad (x, y) \in \mathbb{R}^2,$$

$a, b, c, d, e, f \in \mathbb{R}$ with $bf - ce \neq 0$. Namely, let $L_\beta : \mathbb{R}[x, y]_{\leq 2k} \rightarrow \mathbb{R}$ be a **Riesz functional** of the sequence β defined by

$$L_\beta(p) := \sum_{\substack{i,j \in \mathbb{Z}_+, \\ 0 \leq i+j \leq 2k}} a_{i,j} \beta_{i,j}, \quad \text{where } p = \sum_{\substack{i,j \in \mathbb{Z}_+, \\ 0 \leq i+j \leq 2k}} a_{i,j} x^i y^j.$$

We define $\tilde{\beta} = \{\tilde{\beta}_{i,j}\}_{i,j \in \mathbb{Z}_+, 0 \leq i+j \leq 2k}$ by

$$\tilde{\beta}_{i,j} = L_\beta(\phi_1(x, y)^i \cdot \phi_2(x, y)^j).$$

By [CF04, Proposition 1.9], β admits a (r -atomic) K -rm if and only if $\tilde{\beta}$ admits a (r -atomic) $\phi(K)$ -rm. We write $\tilde{\beta} = \phi(\beta)$ and $\mathcal{M}(k; \tilde{\beta}) = \phi(\mathcal{M}(k; \beta))$.

2.3. Generalized Schur complements. Let

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \mathbb{R}^{(n+m) \times (n+m)}$$

be a real matrix where $A \in \mathbb{R}^{n \times n}$, $B \in \mathbb{R}^{n \times m}$, $C \in \mathbb{R}^{m \times n}$ and $D \in \mathbb{R}^{m \times m}$. The **generalized Schur complement** [Zha05] of A (resp. D) in M is defined by

$$M/A = D - CA^\dagger B \quad (\text{resp. } M/D = A - BD^\dagger C),$$

where A^\dagger (resp. D^\dagger) stands for the Moore–Penrose inverse of A (resp. D).

The following lemma will be frequently used in the proofs.

Lemma 2.1. *Let $n, m \in \mathbb{N}$ and*

$$M = \begin{pmatrix} A & B \\ B^T & C \end{pmatrix} \in S_{n+m},$$

where $A \in S_n$, $B \in \mathbb{R}^{n \times m}$ and $C \in S_m$. If $\text{rank } M = \text{rank } A$, then the matrix equation

$$(2.2) \quad \begin{pmatrix} A \\ B^T \end{pmatrix} W = \begin{pmatrix} B \\ C \end{pmatrix},$$

where $W \in \mathbb{R}^{n \times m}$, is solvable and the solutions are precisely the solutions of the matrix equation $AW = B$. In particular, $W = A^\dagger B$ satisfies (2.2).

Proof. The assumption $\text{rank } M = \text{rank } A$ implies that

$$(2.3) \quad \begin{pmatrix} A \\ B^T \end{pmatrix} W = \begin{pmatrix} AW \\ B^T W \end{pmatrix} = \begin{pmatrix} B \\ C \end{pmatrix}$$

for some $W \in \mathbb{R}^{n \times m}$. So the equation (2.2) is solvable. In particular, $AW = B$. It remains to prove that any solution W to $AW = B$ is also a solution to (2.3). Note that all the solutions of the equation $A\widetilde{W} = B$ are

$$(2.4) \quad \widetilde{W} = A^\dagger B + Z,$$

where each column of $Z \in \mathbb{R}^{n \times m}$ is an arbitrary vector from $\ker A$. So W satisfying (2.3) is also of the form $A^\dagger B + Z_0$ for some $Z_0 \in \mathbb{R}^{n \times m}$ with columns belonging to $\ker A$. We have that

$$(2.5) \quad C = B^T W = B^T (A^\dagger B + Z_0) = B^T A^\dagger B + B^T Z_0 = B^T A^\dagger B,$$

where we used the fact that each column of B belongs to $\mathcal{C}(A)$ and $\ker(A)^\perp = \mathcal{C}(A)$. Replacing W with any \widetilde{W} of the form (2.4) in the calculation (2.5) gives the same result, which proves the statement of the proposition. \square

The following theorem is a characterization of psd 2×2 block matrices.

Theorem 2.2 ([Alb69]). *Let*

$$M = \begin{pmatrix} A & B \\ B^T & C \end{pmatrix} \in S_{n+m}$$

be a real symmetric matrix where $A \in S_n$, $B \in \mathbb{R}^{n \times m}$ and $C \in S_m$. Then:

- (1) *The following conditions are equivalent:*
 - (a) $M \succeq 0$.
 - (b) $C \succeq 0$, $\mathcal{C}(B^T) \subseteq \mathcal{C}(C)$ and $M/C \succeq 0$.
 - (c) $A \succeq 0$, $\mathcal{C}(B) \subseteq \mathcal{C}(A)$ and $M/A \succeq 0$.
- (2) *If $M \succeq 0$, then*

$$\text{rank } M = \text{rank } A + \text{rank } M/A = \text{rank } C + \text{rank } M/C.$$

2.4. Extension principle.

Proposition 2.3. *Let $\mathcal{A} \in S_n$ be positive semidefinite, Q a subset of the set $\{1, \dots, n\}$ and $\mathcal{A}|_Q$ the restriction of \mathcal{A} to the rows and columns from the set Q . If $\mathcal{A}|_Q v = 0$ for a nonzero vector v , then $\mathcal{A}\widehat{v} = 0$, where \widehat{v} is a vector with the only nonzero entries in the rows from Q and such that the restriction $\widehat{v}|_Q$ to the rows from Q equals to v .*

Proof. See [Fia95, Proposition 2.4] or [Zal22a, Lemma 2.4] for an alternative proof. \square

2.5. Partially positive semidefinite matrices and their completions. A **partial matrix** $A = (a_{i,j})_{i,j=1}^n$ is a matrix of real numbers $a_{i,j} \in \mathbb{R}$, where some of the entries are not specified.

A partial symmetric matrix $A = (a_{i,j})_{i,j=1}^n$ is **partially positive semidefinite (ppsd)** (resp. **partially positive definite (ppd)**) if the following two conditions hold:

- (1) $a_{i,j}$ is specified if and only if $a_{j,i}$ is specified and $a_{i,j} = a_{j,i}$.
- (2) All fully specified principal minors of A are psd (resp. pd).

For $n \in \mathbb{N}$ write $[n] := \{1, 2, \dots, n\}$. We denote by A_{Q_1, Q_2} the submatrix of $A \in \mathbb{R}^{n \times n}$ consisting of the rows indexed by the elements of $Q_1 \subseteq [n]$ and the columns indexed by the elements of $Q_2 \subseteq [n]$. In case $Q := Q_1 = Q_2$, we write $A_Q := A_{Q, Q}$ for short.

It is well-known that a ppsd matrix $A(\mathbf{x})$ of the form as in Lemma 2.4 below admits a psd completion (This follows from the fact that the corresponding graph is chordal, see e.g., [GJSW84, Dan92, BW11]). Since we will need an additional information about the rank of the completion $A(x_0)$ and the explicit interval of all possible x_0 for our results, we give a proof of Lemma 2.4 based on the use of generalized Schur complements.

Lemma 2.4. *Let $A(\mathbf{x})$ be a partially positive semidefinite symmetric matrix of size $n \times n$ with the missing entries in the positions (i, j) and (j, i) , $1 \leq i < j \leq n$. Let*

$$A_1 = (A(\mathbf{x}))_{[n] \setminus \{i, j\}}, \quad a = (A(\mathbf{x}))_{[n] \setminus \{i, j\}, \{i\}}, \quad b = (A(\mathbf{x}))_{[n] \setminus \{i, j\}, \{j\}}, \quad \alpha = (A(\mathbf{x}))_{i, i}, \quad \gamma = (A(\mathbf{x}))_{j, j}.$$

Let

$$A_2 = (A(\mathbf{x}))_{[n] \setminus \{j\}} = \begin{pmatrix} A_1 & a \\ a^T & \alpha \end{pmatrix} \in S_{n-1}, \quad A_3 = (A(\mathbf{x}))_{[n] \setminus \{i\}} = \begin{pmatrix} A_1 & b \\ b^T & \gamma \end{pmatrix} \in S_{n-1},$$

and

$$x_{\pm} := b^T A_1^\dagger a \pm \sqrt{(A_2/A_1)(A_3/A_1)} \in \mathbb{R}.$$

Then:

- (i) $A(x_0)$ is positive semidefinite if and only if $x_0 \in [x_-, x_+]$.
- (ii)

$$\text{rank } A(x_0) = \begin{cases} \max \{ \text{rank } A_2, \text{rank } A_3 \}, & \text{for } x_0 \in \{x_-, x_+\}, \\ \max \{ \text{rank } A_2, \text{rank } A_3 \} + 1, & \text{for } x_0 \in (x_-, x_+). \end{cases}$$

(iii) The following statements are equivalent:

- (a) $x_- = x_+$.
- (b) $A_2/A_1 = 0$ or $A_3/A_1 = 0$.
- (c) $\text{rank } A_2 = \text{rank } A_1$ or $\text{rank } A_3 = \text{rank } A_1$.

Proof. We write

$$A(\mathbf{x}) = \begin{pmatrix} A_{11} & a_{12} & A_{13} & a_{14} & A_{15} \\ (a_{12})^T & \alpha & (a_{23})^T & \mathbf{x} & (a_{25})^T \\ (A_{13})^T & a_{23} & A_{33} & a_{34} & a_{35} \\ (a_{14})^T & \mathbf{x} & (a_{34})^T & \gamma & (a_{45})^T \\ (A_{15})^T & a_{25} & (A_{35})^T & a_{45} & A_{55} \end{pmatrix} \in \begin{pmatrix} S_{i-1} & \mathbb{R}^{(i-1) \times 1} & \mathbb{R}^{(i-1) \times (j-i-1)} & \mathbb{R}^{(i-1) \times 1} & \mathbb{R}^{(i-1) \times (n-j)} \\ \mathbb{R}^{1 \times (i-1)} & \mathbb{R} & \mathbb{R}^{1 \times (j-i-1)} & \mathbb{R} & \mathbb{R}^{1 \times (n-j)} \\ \mathbb{R}^{(j-i-1) \times (i-1)} & \mathbb{R}^{(j-i-1) \times 1} & S_{j-i-1} & \mathbb{R}^{(j-i-1) \times 1} & \mathbb{R}^{(j-i-1) \times (n-j)} \\ \mathbb{R}^{1 \times (i-1)} & \mathbb{R} & \mathbb{R}^{1 \times (j-i-1)} & \mathbb{R} & \mathbb{R}^{1 \times (n-j)} \\ \mathbb{R}^{(n-j) \times (i-1)} & \mathbb{R}^{(n-j) \times 1} & \mathbb{R}^{(n-j) \times (j-i-1)} & \mathbb{R}^{(n-j) \times 1} & S_{n-j} \end{pmatrix}$$

Let P be a permutation matrix, which changes the order of columns to

$$1, 2, \dots, i-1, i+1, \dots, j-1, j+1, \dots, n, i, j.$$

Then

$$P^T A(\mathbf{x})P = \begin{pmatrix} A_{11} & A_{13} & A_{15} & a_{12} & a_{14} \\ (A_{13})^T & A_{33} & A_{35} & a_{23} & a_{34} \\ (A_{15})^T & (A_{35})^T & A_{55} & a_{25} & a_{45} \\ (a_{12})^T & (a_{23})^T & (a_{25})^T & \alpha & \mathbf{x} \\ (a_{14})^T & (a_{34})^T & (a_{45})^T & \mathbf{x} & \gamma \end{pmatrix}$$

Note that

$$(2.6) \quad P^T A(\mathbf{x})P = \begin{pmatrix} A_1 & a & b \\ a^T & \alpha & \mathbf{x} \\ b^T & \mathbf{x} & \gamma \end{pmatrix} \quad \text{and} \quad P^T A(\mathbf{x})P \succeq 0 \Leftrightarrow A(\mathbf{x}) \succeq 0.$$

Lemma 2.4 with the missing entries in the positions $(n-1, n)$ and $(n, n-1)$ was proved in [Zal21, Lemma 2.11] using computations with generalized Schur complements under one additional assumption:

$$(2.7) \quad A_1 \text{ is invertible} \quad \text{or} \quad \text{rank } A_1 = \text{rank } A_2.$$

Here we explain why the assumption (2.7) can be removed from [Zal21, Lemma 2.11]. The proof of [Zal21, Lemma 2.11] is separated into two cases: $A_2/A_1 > 0$ and $A_2/A_1 = 0$. The case $A_2/A_1 = 0$ does not use (2.7). Assume now that $A_2/A_1 > 0$ or equivalently $\text{rank } A_2 > \text{rank } A_1$. Invertibility of A_1 (and by $A_2/A_1 > 0$ also A_2 is invertible) is used in the proof of [Zal21, Lemma 2.11] for the application of the quotient formula ([CH69])

$$(2.8) \quad (A(x)/A_2) = (A(x)/A_1)/(A_2/A_1),$$

where

$$A(x)/A_1 = \begin{pmatrix} A_2/A_1 & \begin{pmatrix} A_1 & b \\ a^T & x \end{pmatrix}/A_1 \\ \begin{pmatrix} A_1 & a \\ b^T & x \end{pmatrix}/A_1 & A_3/A_1 \end{pmatrix}$$

However, the formula (2.8) has been generalized [CHM74, Theorem 4] to noninvertible A_1, A_2 , where all Schur complements are the generalized ones, under the conditions:

$$(2.9) \quad (b \ x)^T \in \mathcal{C}(A_2) \quad \text{and} \quad a \in \mathcal{C}(A_1).$$

So if we show that the conditions (2.9) hold, the same proof as in [Zal21, Lemma 2.11] can be applied in the case A_1 is singular. From A_2 (resp. A_3) being psd, $a \in \mathcal{C}(A_1)$ (resp. $b \in \mathcal{C}(A_1)$) follows by Theorem 2.2, used for $(M, A) := (A_2, A_1)$ (resp. $(M, A) := (A_3, A_1)$). The assumption $A_2/A_1 > 0$ implies that $(a \ \alpha)^T \notin \mathcal{C}((A_1 \ a^T)^T)$. Since $a \in \mathcal{C}(A_1)$, it follows that $(0 \ 1)^T \in \mathcal{C}(A_2)$. Hence, $(b \ x)^T \in \mathcal{C}(A_2)$ for every $x \in \mathbb{R}$, which concludes the proof of (2.9). \square

2.6. Hamburger TMP. Let $k \in \mathbb{N}$. For $v = (v_0, \dots, v_{2k}) \in \mathbb{R}^{2k+1}$ we define the corresponding Hankel matrix as

$$(2.10) \quad A_v := (v_{i+j})_{i,j=0}^k = \begin{pmatrix} v_0 & v_1 & v_2 & \cdots & v_k \\ v_1 & v_2 & \ddots & \ddots & v_{k+1} \\ v_2 & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & v_{2k-1} \\ v_k & v_{k+1} & \cdots & v_{2k-1} & v_{2k} \end{pmatrix} \in S_{k+1}.$$

We denote by $\mathbf{v}_j := (v_{j+\ell})_{\ell=0}^k$ the $(j+1)$ -th column of A_v , $0 \leq j \leq k$, i.e.,

$$A_v = \begin{pmatrix} \mathbf{v}_0 & \cdots & \mathbf{v}_k \end{pmatrix}.$$

As in [CF91], the **rank** of v , denoted by $\text{rank } v$, is defined by

$$\text{rank } v = \begin{cases} k+1, & \text{if } A_v \text{ is nonsingular,} \\ \min \{i: \mathbf{v}_i \in \text{span}\{\mathbf{v}_0, \dots, \mathbf{v}_{i-1}\}\}, & \text{if } A_v \text{ is singular.} \end{cases}$$

For $m \leq k$ we denote the upper left-hand corner $(v_{i+j})_{i,j=0}^m \in S_{m+1}$ of A_v of size $m+1$ by $A_v(m)$. A sequence v is called **positively recursively generated (prg)** if for $r = \text{rank } v$ the following two conditions hold:

- $A_v(r-1) \succ 0$.
- If $r < k+1$, denoting

$$(2.11) \quad (\varphi_0, \dots, \varphi_{r-1}) := A_v(r-1)^{-1}(v_r, \dots, v_{2r-1})^T,$$

the equality

$$(2.12) \quad v_j = \varphi_0 v_{j-r} + \cdots + \varphi_{r-1} v_{j-1}$$

holds for $j = r, \dots, 2k$.

The solution to the \mathbb{R} -TMP is the following.

Theorem 2.5 ([CF91, Theorems 3.9–3.10]). *For $k \in \mathbb{N}$ and $v = (v_0, \dots, v_{2k}) \in \mathbb{R}^{2k+1}$ with $v_0 > 0$, the following statements are equivalent:*

- (1) *There exists a \mathbb{R} -representing measure for β .*
- (2) *There exists a $(\text{rank } A_v)$ -atomic \mathbb{R} -representing measure for β .*
- (3) *A_v is positive semidefinite and one of the following holds:*
 - (a) *$A_v(k-1)$ is positive definite.*
 - (b) *$\text{rank } A_v(k-1) = \text{rank } A_v$.*
- (4) *v is positively recursively generated.*

2.7. TMP on the unit circle. The solution to the $\mathcal{Z}(x^2 + y^2 - 1)$ -TMP is the following.

Theorem 2.6 ([CF02, Theorem 2.1]). *Let $p(x, y) = x^2 + y^2 - 1$ and $\beta := \beta^{(2k)} = (\beta_{i,j})_{i,j \in \mathbb{Z}_+, i+j \leq 2k}$, where $k \geq 2$. Then the following statements are equivalent:*

- (1) *β has a $\mathcal{Z}(p)$ -representing measure.*
- (2) *β has a $(\text{rank } \mathcal{M}(k))$ -atomic $\mathcal{Z}(p)$ -representing measure.*
- (3) *$\mathcal{M}(k)$ is positive semidefinite and the relations $\beta_{2+i,j} + \beta_{i,2+j} = \beta_{i,j}$ hold for every $i, j \in \mathbb{Z}_+$ with $i+j \leq 2k-2$.*

2.8. Parabolic TMP. We will need the following solution to the parabolic TMP (see [Zal23, Theorem 3.7]).

Theorem 2.7. *Let $p(x, y) = x - y^2$ and $\beta := \beta^{(2k)} = (\beta_{i,j})_{i,j \in \mathbb{Z}_+, i+j \leq 2k}$, where $k \geq 2$. Let*

$$\mathcal{B} = \{1, Y, X, XY, X^2, X^2Y, \dots, X^i, X^iY, \dots, X^{k-1}, X^{k-1}Y, X^k\}.$$

Then the following statements are equivalent:

- (1) *β has a $\mathcal{Z}(p)$ -representing measure.*
- (2) *β has a $(\text{rank } \mathcal{M}(k))$ -atomic $\mathcal{Z}(p)$ -representing measure.*

- (3) $\mathcal{M}(k)$ is positive semidefinite, the relations $\beta_{1+i,j} = \beta_{i,2+j}$ hold for every $i, j \in \mathbb{Z}_+$ with $i + j \leq 2k - 2$ and one of the following statements holds:
- (a) $(\mathcal{M}(k))_{\mathcal{B} \setminus \{X^k\}}$ is positive definite.
 - (b) $\text{rank}(\mathcal{M}(k))_{\mathcal{B} \setminus \{X^k\}} = \text{rank} \mathcal{M}(k)$.
- (4) The relations $\beta_{1+i,j} = \beta_{i,2+j}$ hold for every $i, j \in \mathbb{Z}_+$ with $i + j \leq 2k - 2$ and $\gamma = (\gamma_0, \dots, \gamma_{4k})$, defined by $\gamma_i = \beta_{\lfloor \frac{i}{2} \rfloor, i \bmod 2}$, admits a \mathbb{R} -representing measure.

Remark 2.8. The equivalence (3) \Leftrightarrow (4) is part of the proof of [Zal23, Theorem 3.7].

3. TMP ON REDUCIBLE CUBICS - CASE REDUCTION

In this section we show that to solve the TMP on reducible cubic curves it suffices, after applying an affine linear transformation, to solve the TMP on 8 canonical forms of curves.

Proposition 3.1. *Let $k \in \mathbb{R}$ and $\beta := \beta^{(2k)} = (\beta_{i,j})_{i,j \in \mathbb{Z}_+, i+j \leq 2k}$. Assume $\mathcal{M}(k; \beta)$ does not satisfy any nontrivial column relation between columns indexed by monomials of degree at most 2, but it satisfies a column relation $p(X, Y) = \mathbf{0}$, where $p \in \mathbb{R}[x, y]$ is a reducible polynomial with $\deg p = 3$. If β admits a representing measure, then there exists an invertible affine linear transformation ϕ of the form (2.1) such that the moment matrix $\phi(\mathcal{M}(k; \beta))$ satisfies a column relation $q(x, y) = 0$, where q has one of the following forms:*

Parallel lines type: $q(x, y) = y(a + y)(b + y)$, $a, b \in \mathbb{R} \setminus \{0\}$, $a \neq b$.

Circular type: $q(x, y) = y(ay + x^2 + y^2)$, $a \in \mathbb{R} \setminus \{0\}$.

Parabolic type: $q(x, y) = y(x - y^2)$.

Hyperbolic type 1: $q(x, y) = y(1 - xy)$.

Hyperbolic type 2: $q(x, y) = y(x + y + axy)$, $a \in \mathbb{R} \setminus \{0\}$.

Hyperbolic type 3: $q(x, y) = y(ay + x^2 - y^2)$, $a \in \mathbb{R}$.

Intersecting lines type: $q(x, y) = yx(y + 1)$,

Mixed type: $q(x, y) = y(1 + ay + bx^2 + cy^2)$, $a, b, c \in \mathbb{R}$, $b \neq 0$.

Remark 3.2. The name of the types of the form q in Proposition 3.1 comes from the type of the conic $\frac{q(x,y)}{y} = 0$. The conic $x + y + axy = 0$, $a \in \mathbb{R} \setminus \{0\}$, is a hyperbola, since the discriminant a^2 is positive. Similarly, the conic $ay + x^2 - y^2 = 0$, $a \in \mathbb{R}$, is a hyperbola, since its discriminant is equal to 4. Clearly, the conic $ay + x^2 + y^2 = 0$, $a \in \mathbb{R}$, is a circle with the center $(0, -\frac{a}{2})$ and radius $\frac{a}{2}$.

Now we prove Proposition 3.1.

Proof of Proposition 3.1. Since $p(x, y)$ is reducible, it is of the form $p = p_1 p_2$, where

$$p_1(x, y) = a_0 + a_1 x + a_2 y \quad \text{with } a_i \in \mathbb{R}, (a_1, a_2) \neq (0, 0),$$

$$p_2(x, y) = b_0 + b_1 x + b_2 y + b_3 x^2 + b_4 xy + b_5 y^2 \quad \text{with } b_i \in \mathbb{R}, (b_3, b_4, b_5) \neq (0, 0, 0).$$

Without loss of generality we can assume that $a_2 \neq 0$, since otherwise we apply the alt $(x, y) \mapsto (y, x)$ to exchange the roles of x and y . Since $a_2 \neq 0$, the alt

$$\phi_1(x, y) = (x, a_0 + a_1 x + a_2 y)$$

is invertible and hence:

$$(3.1) \quad \text{A sequence } \phi_1(\beta) \text{ has a moment matrix } \phi_1(\mathcal{M}(k; \beta)) \text{ satisfying the column relation} \\ c_0Y + c_1X + c_2Y^2 + c_3X^2Y + c_4XY^2 + c_5Y^3 = \mathbf{0} \text{ with } c_i \in \mathbb{R}, (c_3, c_4, c_5) \neq (0, 0, 0).$$

We separate two cases according to the value of c_3 .

Case 1: $c_3 = 0$. In this case (3.1) is equal to

$$(3.2) \quad \text{A sequence } \phi_1(\beta) \text{ has a moment matrix } \phi_1(\mathcal{M}(k; \beta)) \text{ satisfying the column relation} \\ c_0Y + c_1XY + c_2Y^2 + c_4XY^2 + c_5Y^3 = \mathbf{0} \text{ with } c_i \in \mathbb{R}, (c_4, c_5) \neq (0, 0).$$

If $c_0 = c_1 = c_2 = 0$, then (3.2) is equal to $c_4XY^2 + c_5Y^3 = \mathbf{0}$. Since by assumption β and hence $\phi_1(\beta)$ admit a rm, supported on

$$\mathcal{Z}(y^2(c_4x + c_5y)) = \mathcal{Z}(y(c_4x + c_5y)),$$

it follows by [CF96] that $c_4XY + c_5Y^2 = \mathbf{0}$ is a nontrivial column relation in $\phi_1(\mathcal{M}(k; \beta))$. Hence, also $\mathcal{M}(k; \beta)$ satisfies a nontrivial column relation between columns indexed by monomials of degree at most 2, which is a contradiction with the assumption of the proposition. Therefore $(c_0, c_1, c_2) \neq (0, 0, 0)$.

Case 1.1: $c_0 \neq 0$. Dividing the relation in (3.2) by c_0 , we get:

$$(3.3) \quad \text{A sequence } \phi_1(\beta) \text{ has a moment matrix } \phi_1(\mathcal{M}(k; \beta)) \text{ satisfying the column relation} \\ Y + \tilde{c}_1XY + \tilde{c}_2Y^2 + \tilde{c}_4XY^2 + \tilde{c}_5Y^3 = \mathbf{0} \text{ with } \tilde{c}_i \in \mathbb{R}, (\tilde{c}_4, \tilde{c}_5) \neq (0, 0).$$

Case 1.1.1: $\tilde{c}_1 = 0$. In this case (3.3) is equivalent to:

$$(3.4) \quad \text{A sequence } \phi_1(\beta) \text{ has a moment matrix } \phi_1(\mathcal{M}(k; \beta)) \text{ satisfying the column relation} \\ Y + \tilde{c}_2Y^2 + \tilde{c}_4XY^2 + \tilde{c}_5Y^3 = \mathbf{0} \text{ with } \tilde{c}_i \in \mathbb{R}, (\tilde{c}_4, \tilde{c}_5) \neq (0, 0).$$

Case 1.1.1.1: $\tilde{c}_4 = 0$. In this case (3.4) is equivalent to

$$(3.5) \quad \text{A sequence } \phi_1(\beta) \text{ has a moment matrix } \phi_1(\mathcal{M}(k; \beta)) \text{ satisfying the column relation} \\ Y + \tilde{c}_2Y^2 + \tilde{c}_5Y^3 = \mathbf{0} \text{ with } \tilde{c}_2 \in \mathbb{R}, \tilde{c}_5 \in \mathbb{R} \setminus \{0\}.$$

The quadratic equation $1 + \tilde{c}_2y + \tilde{c}_5y^2 = 0$ must have two different real nonzero solutions, otherwise $\mathcal{Z}(y(1 + \tilde{c}_2x + \tilde{c}_5y))$ is a union of two parallel lines. Then it follows by [CF96] that there is a nontrivial column relation in $\mathcal{M}(k; \beta)$ between columns indexed by monomials of degree at most 2, which is a contradiction with the assumption of the proposition. So we have the parallel lines type relation from the proposition.

Case 1.1.1.2: $\tilde{c}_4 \neq 0$. In this case the alt

$$\phi_2(x, y) = (-\tilde{c}_2 - \tilde{c}_4x - \tilde{c}_5y, y)$$

is invertible and applying it to $\phi_1(\beta)$, we obtain:

A sequence $(\phi_2 \circ \phi_1)(\beta)$ has a moment matrix $(\phi_2 \circ \phi_1)(\mathcal{M}(k; \beta))$ satisfying the hyperbolic type 1 relation from the proposition.

Case 1.1.2: $\tilde{c}_1 \neq 0$. We apply the alt

$$\phi_3(x, y) = (1 + \tilde{c}_1x, y)$$

to $\phi_1(\beta)$ and obtain:

(3.6) A sequence $(\phi_3 \circ \phi_1)(\beta)$ has a moment matrix $(\phi_3 \circ \phi_1)(\mathcal{M}(k; \beta))$ satisfying the column relation $XY + \widehat{c}_2 Y^2 + \widehat{c}_4 XY^2 + \widehat{c}_5 Y^3 = \mathbf{0}$ with $\widehat{c}_i \in \mathbb{R}$, $(\widehat{c}_4, \widehat{c}_5) \neq (0, 0)$.

Case 1.1.2.1: $\widehat{c}_4 \neq 0$. We apply the alt

$$\phi_4(x, y) = \left(x - \frac{\widehat{c}_5}{\widehat{c}_4} y, y \right)$$

to $(\phi_3 \circ \phi_1)(\beta)$ and obtain:

(3.7) A sequence $(\phi_4 \circ \phi_3 \circ \phi_1)(\beta)$ has a moment matrix $(\phi_4 \circ \phi_3 \circ \phi_1)(\mathcal{M}(k; \beta))$ satisfying the column relation $XY + \check{c}_2 Y^2 + \widehat{c}_4 XY^2 = \mathbf{0}$ with $\check{c}_2, \widehat{c}_4 \in \mathbb{R}$, $\widehat{c}_4 \neq 0$.

Case 1.1.2.1.1: $\check{c}_2 = 0$. In this case the relation in (3.7) is of the form

$$XY + \widehat{c}_4 XY^2 = \mathbf{0} \quad \text{with } \widehat{c}_4 \in \mathbb{R} \setminus \{0\}.$$

Applying the alt

$$\phi_5(x, y) = (x, \widehat{c}_4 y)$$

to $(\phi_4 \circ \phi_3 \circ \phi_1)(\beta)$ we obtain:

A sequence $(\phi_5 \circ \phi_4 \circ \phi_3 \circ \phi_1)(\beta)$ has a moment matrix $(\phi_5 \circ \phi_4 \circ \phi_3 \circ \phi_1)(\mathcal{M}(k; \beta))$ satisfying the intersecting lines type relation from the proposition.

Case 1.1.2.1.2: $\check{c}_2 \neq 0$. We apply the alt

$$\phi_6(x, y) = (x, \check{c}_2 y)$$

to $(\phi_4 \circ \phi_3 \circ \phi_1)(\beta)$ and obtain:

A sequence $(\phi_6 \circ \phi_4 \circ \phi_3 \circ \phi_1)(\beta)$ has a moment matrix $(\phi_6 \circ \phi_4 \circ \phi_3 \circ \phi_1)(\mathcal{M}(k; \beta))$ satisfying the hyperbolic type 2 relation in the proposition.

Case 1.1.2.2: $\widehat{c}_4 = 0$. In this case (3.6) is equivalent to:

(3.8) A sequence $(\phi_3 \circ \phi_1)(\beta)$ has a moment matrix $(\phi_3 \circ \phi_1)(\mathcal{M}(k; \beta))$ satisfying the column relation $XY + \widehat{c}_2 Y^2 + \widehat{c}_5 Y^3 = \mathbf{0}$ with $\widehat{c}_2, \widehat{c}_5 \in \mathbb{R}$, $\widehat{c}_5 \neq 0$.

Case 1.1.2.2.1: $\widetilde{c}_2 = 0$. Applying the alt

$$\phi_7(x, y) = (x, -\widehat{c}_5 y),$$

to $(\phi_3 \circ \phi_1)(\beta)$ we obtain:

A sequence $(\phi_7 \circ \phi_3 \circ \phi_1)(\beta)$ has a moment matrix $(\phi_7 \circ \phi_3 \circ \phi_1)(\mathcal{M}(k; \beta))$ satisfying the parabolic type relation in the proposition.

Case 1.1.2.2.2: $\widetilde{c}_2 \neq 0$. Applying the alt

$$\phi_8(x, y) = (x, \widehat{c}_2 y)$$

to $(\phi_3 \circ \phi_1)(\beta)$ and obtain:

(3.9) A sequence $(\phi_8 \circ \phi_3 \circ \phi_1)(\beta)$ has a moment matrix $(\phi_8 \circ \phi_3 \circ \phi_1)(\mathcal{M}(k; \beta))$ satisfying the column relation $XY + Y^2 + \check{c}_5 Y^3 = \mathbf{0}$ with $\check{c}_5 \in \mathbb{R}$, $\check{c}_5 \neq 0$.

Further on, the relation in (3.9) is equivalent to

$$(3.10) \quad (\check{c}_5)^{-1}(XY + Y^2) + Y^3 = \mathbf{0} \quad \text{with } \check{c}_5 \in \mathbb{R}, \check{c}_5 \neq 0.$$

Finally, applying the alt

$$\phi_9(x, y) = ((-\check{c}_5)^{-1}(x + y), y)$$

to $(\phi_8 \circ \phi_3 \circ \phi_1)(\beta)$, we obtain:

A sequence $(\phi_9 \circ \phi_8 \circ \phi_3 \circ \phi_1)(\beta)$ has a moment matrix $(\phi_9 \circ \phi_8 \circ \phi_3 \circ \phi_1)(\mathcal{M}(k; \beta))$ satisfying the parabolic type relation in the proposition.

Case 1.2: $c_0 = 0$. In this case (3.2) is equivalent to:

$$(3.11) \quad \text{A sequence } \phi_1(\beta) \text{ has a moment matrix } \phi_1(\mathcal{M}(k; \beta)) \text{ satisfying the column relation} \\ c_1XY + c_2Y^2 + c_4XY^2 + c_5Y^3 = \mathbf{0} \text{ with } c_i \in \mathbb{R}, (c_4, c_5) \neq (0, 0).$$

Assume that $c_1 = 0$. Since by assumption β and hence $\phi_1(\beta)$ admits a rm, supported on

$$\mathcal{Z}(y^2(c_2 + c_4x + c_5y)) = \mathcal{Z}(y(c_2 + c_4x + c_5y)),$$

it follows by [CF96] that $c_2Y + c_4XY + c_5Y^2 = \mathbf{0}$ is a nontrivial column relation in $\phi_1(\mathcal{M}(k; \beta))$. Hence, also $\mathcal{M}(k; \beta)$ satisfies a nontrivial column relation between columns indexed by monomials of degree at most 2, which is a contradiction with the assumption of the proposition. Hence, $c_1 \neq 0$. Applying the alt $(x, y) \mapsto (c_1x, y)$ to $\phi_1(\beta)$, we obtain a sequence with the moment matrix satisfying the column relation of the form (3.6) and we can proceed as in the Case 1.1.2 above.

Case 2: $c_3 \neq 0$. Applying the alt

$$\phi_{10}(x, y) = (\sqrt{|c_3|}x, y)$$

to $\phi_1(\beta)$, we obtain:

$$(3.12) \quad \text{A sequence } (\phi_{10} \circ \phi_1)(\beta) \text{ has a moment matrix } (\phi_{10} \circ \phi_1)(\mathcal{M}(k; \beta)) \text{ satisfying} \\ \text{the column relation } c_0Y + \tilde{c}_1XY + c_2Y^2 + \frac{|c_3|}{c_3}X^2Y + \tilde{c}_4XY^2 + c_5Y^3 = \mathbf{0} \text{ with } c_i, \tilde{c}_i \in \mathbb{R}.$$

Case 2.1: $\tilde{c}_1 = 0$. In this case (3.12) is equivalent to:

$$(3.13) \quad \text{A sequence } (\phi_{10} \circ \phi_1)(\beta) \text{ has a moment matrix } (\phi_{10} \circ \phi_1)(\mathcal{M}(k; \beta)) \text{ satisfying} \\ \text{the column relation } c_0Y + c_2Y^2 + \frac{|c_3|}{c_3}X^2Y + \tilde{c}_4XY^2 + c_5Y^3 = \mathbf{0} \text{ with } c_i, \tilde{c}_i \in \mathbb{R}.$$

Case 2.1.1: $c_0 = 0$. Dividing the relation in (3.13) with $\frac{|c_3|}{c_3}$, (3.13) is equivalent to:

$$(3.14) \quad \text{A sequence } (\phi_{10} \circ \phi_1)(\beta) \text{ has a moment matrix } (\phi_{10} \circ \phi_1)(\mathcal{M}(k; \beta)) \text{ satisfying} \\ \text{the column relation } \tilde{c}_2Y^2 + X^2Y + \hat{c}_4XY^2 + \tilde{c}_5Y^3 = \mathbf{0} \text{ with } \tilde{c}_2, \hat{c}_4, \tilde{c}_5 \in \mathbb{R}.$$

Applying the alt

$$\phi_{11}(x, y) = \left(x + \frac{\hat{c}_4}{2}y, y\right)$$

to $(\phi_{10} \circ \phi_1)(\beta)$, we obtain:

$$(3.15) \quad \text{A sequence } (\phi_{11} \circ \phi_{10} \circ \phi_1)(\beta) \text{ has a moment matrix } (\phi_{11} \circ \phi_{10} \circ \phi_1)(\mathcal{M}(k; \beta)) \\ \text{satisfying the column relation } \check{c}_2Y^2 + X^2Y + \check{c}_5Y^3 = \mathbf{0} \text{ with } \check{c}_2, \check{c}_5 \in \mathbb{R}.$$

Case 2.1.1.1: $\check{c}_5 = 0$. Since by assumption of the proposition, $(\phi_{11} \circ \phi_{10} \circ \phi_1)(\beta)$ admits a rm, supported on $\mathcal{Z}(y(\check{c}_2y + x^2))$, \check{c}_2 in (3.15) cannot be equal to 0. Indeed, $\check{c}_2 = 0$ would imply that $\mathcal{Z}(y(\check{c}_2y + x^2)) = \mathcal{Z}(yx^2) = \mathcal{Z}(yx)$ and by [CF96], $XY = \mathbf{0}$ would be a nontrivial column relation in $(\phi_{11} \circ \phi_{10} \circ \phi_1)(\mathcal{M}(k; \beta))$. Hence, also $\mathcal{M}(k; \beta)$ would satisfy a nontrivial column relation between columns indexed by monomials of degree at most 2, which is a contradiction with the assumption of the proposition. Since $\check{c}_2 \neq 0$, after applying the alt

$$\phi_{12}(x, y) = (x, -\check{c}_2y)$$

to $(\phi_{11} \circ \phi_{10} \circ \phi_1)(\beta)$, we obtain:

A sequence $(\phi_{12} \circ \phi_{11} \circ \phi_{10} \circ \phi_1)(\beta)$ has a moment matrix $(\phi_{12} \circ \phi_{11} \circ \phi_{10} \circ \phi_1)(\mathcal{M}(k; \beta))$ satisfying the parabolic type relation in the proposition.

Case 2.1.1.2: $\check{c}_5 > 0$. Applying the alt

$$\phi_{13}(x, y) = (x, \sqrt{\check{c}_5}y)$$

to $(\phi_{11} \circ \phi_{10} \circ \phi_1)(\beta)$ we obtain:

A sequence $(\phi_{13} \circ \phi_{11} \circ \phi_{10} \circ \phi_1)(\beta)$ has a moment matrix $(\phi_{13} \circ \phi_{11} \circ \phi_{10} \circ \phi_1)(\mathcal{M}(k; \beta))$ satisfying the circular type relation in the proposition.

Case 2.1.1.3: $\check{c}_5 < 0$. Applying the alt

$$\phi_{14}(x, y) = (x, \sqrt{-\check{c}_5}y)$$

to $(\phi_{11} \circ \phi_{10} \circ \phi_1)(\beta)$, we obtain:

A sequence $(\phi_{14} \circ \phi_{11} \circ \phi_{10} \circ \phi_1)(\beta)$ has a moment matrix $(\phi_{14} \circ \phi_{11} \circ \phi_{10} \circ \phi_1)(\mathcal{M}(k; \beta))$ satisfying the hyperbolic type 3 relation in the proposition.

Case 2.1.2: $c_0 \neq 0$. Dividing the relation in (3.13) with c_0 , (3.13) is equivalent to:

(3.16) A sequence $(\phi_{10} \circ \phi_1)(\beta)$ has a moment matrix $(\phi_{10} \circ \phi_1)(\mathcal{M}(k; \beta))$ satisfying the column relation $Y + \tilde{c}_2Y^2 + \tilde{c}_3X^2Y + \hat{c}_4XY^2 + \tilde{c}_5Y^3 = \mathbf{0}$ with $\tilde{c}_i, \hat{c}_4 \in \mathbb{R}, \tilde{c}_3 \neq 0$.

Applying the alt

$$\phi_{15}(x, y) = \left(x + \frac{\hat{c}_4}{2\tilde{c}_3}, y \right)$$

to $(\phi_{10} \circ \phi_1)(\beta)$, we obtain:

A sequence $(\phi_{15} \circ \phi_{10} \circ \phi_1)(\beta)$ has a moment matrix $(\phi_{15} \circ \phi_{10} \circ \phi_1)(\mathcal{M}(k; \beta))$ satisfying the mixed type relation in the proposition.

Case 2.2: $\tilde{c}_1 \neq 0$. Dividing the relation in (3.12) with $\frac{|c_3|}{c_3}$, (3.12) is equivalent to:

(3.17) A sequence $(\phi_{10} \circ \phi_1)(\beta)$ has a moment matrix $(\phi_{10} \circ \phi_1)(\mathcal{M}(k; \beta))$ satisfying the column relation $\hat{c}_0Y + \hat{c}_1XY + \hat{c}_2Y^2 + X^2Y + \hat{c}_4XY^2 + \hat{c}_5Y^3 = \mathbf{0}$ with $\hat{c}_i \in \mathbb{R}, \hat{c}_1 \neq 0$.

Now we apply the alt

$$\phi_{16}(x, y) = \left(x + \frac{\hat{c}_1}{2}, y \right)$$

to $(\phi_{10} \circ \phi_1)(\beta)$ and obtain:

$$(3.18) \quad \begin{aligned} & \text{A sequence } (\phi_{16} \circ \phi_{10} \circ \phi_1)(\beta) \text{ has a moment matrix } (\phi_{16} \circ \phi_{10} \circ \phi_1)(\mathcal{M}(k; \beta)) \\ & \text{satisfying the column relation } \check{c}_0 Y + \check{c}_2 Y^2 + X^2 Y + \check{c}_4 X Y^2 + \check{c}_5 Y^3 = \mathbf{0} \text{ with } \check{c}_i \in \mathbb{R}. \end{aligned}$$

Case 2.2.1: $\check{c}_0 = 0$. In this case the relation in (3.18) becomes equal to the relation in (3.14) from the Case 2.1.1, so we can proceed as above.

Case 2.2.2: $\check{c}_0 \neq 0$. Dividing the relation in (3.18) with \check{c}_0 , it becomes equal to the relation in (3.16) from the Case 2.1.2, so we can proceed as above. \square

4. SOLVING THE TMP ON CANONICAL REDUCIBLE CUBIC CURVES

Let $\beta = \{\beta_i\}_{i \in \mathbb{Z}_+^2, |i| \leq 2k}$ be a sequence of degree $2k$, $k \in \mathbb{N}$, and

$$(4.1) \quad \mathcal{C} = \{1, X, Y, X^2, XY, Y^2, \dots, X^k, X^{k-1}Y, \dots, Y^k\}$$

the set of rows and columns of the moment matrix $\mathcal{M}(k; \beta)$ in the degree-lexicographic order. Let

$$(4.2) \quad p(x, y) = y \cdot c(x, y) \in \mathbb{R}[x, y]_{\leq 3}$$

be a polynomial of degree 3 in one of the canonical forms from Proposition 3.1. Hence, $c(x, y)$ a polynomial of degree 2. β will have a $\mathcal{Z}(p)$ -rm if and only if it can be decomposed as

$$(4.3) \quad \beta = \beta^{(\ell)} + \beta^{(c)},$$

where

$$\begin{aligned} \beta^{(\ell)} &:= \{\beta_i^{(\ell)}\}_{i \in \mathbb{Z}_+^2, |i| \leq 2k} \quad \text{has a representing measure on } y = 0, \\ \beta^{(c)} &:= \{\beta_i^{(c)}\}_{i \in \mathbb{Z}_+^2, |i| \leq 2k} \quad \text{has a representing measure on the conic } c(x, y) = 0, \end{aligned}$$

and the sum in (4.3) is a component-wise sum. On the level of moment matrices, (4.3) is equivalent to

$$(4.4) \quad \mathcal{M}(k; \beta) = \mathcal{M}(k; \beta^{(\ell)}) + \mathcal{M}(k; \beta^{(c)}).$$

Note that if β has a $\mathcal{Z}(p)$ -rm, then the matrix $\mathcal{M}(k; \beta)$ satisfies the relation $p(X, Y) = \mathbf{0}$ and it must be rg, i.e.,

$$(4.5) \quad X^i Y^j p(X, Y) = \mathbf{0} \quad \text{for } i, j = 0, \dots, k-3 \text{ such that } i+j \leq k-3.$$

We write $\vec{X}^{(0,k)} := (1, X, \dots, X^k)$. Let $\mathcal{T} \subseteq \mathcal{C}$ be a subset, such that the columns from \mathcal{T} span the column space $\mathcal{C}(\mathcal{M}(k; \beta))$ and

$$(4.6) \quad \begin{aligned} & P \text{ is a permutation matrix such that moment matrix } \widetilde{\mathcal{M}}(k; \beta) := P\mathcal{M}(k; \beta)P^T \\ & \text{has rows and columns indexed in the order } \vec{X}^{(0,k)}, \mathcal{T} \setminus \vec{X}^{(0,k)}, \mathcal{C} \setminus (\vec{X}^{(0,k)} \cup \mathcal{T}). \end{aligned}$$

In this new order of rows and columns, (4.4) becomes equivalent to

$$(4.7) \quad \widetilde{\mathcal{M}}(k; \beta) = \widetilde{\mathcal{M}}(k; \beta^{(\ell)}) + \widetilde{\mathcal{M}}(k; \beta^{(c)}).$$

We write

$$(4.8) \quad \widetilde{\mathcal{M}}(k; \beta) = \begin{pmatrix} (\vec{X}^{(0,k)})^T \\ (\mathcal{T} \setminus \vec{X}^{(0,k)})^T \\ (\mathcal{C} \setminus (\vec{X}^{(0,k)} \cup \mathcal{T}))^T \end{pmatrix} \begin{pmatrix} \vec{X}^{(0,k)} & \mathcal{T} \setminus \vec{X}^{(0,k)} & \mathcal{C} \setminus (\vec{X}^{(0,k)} \cup \mathcal{T}) \\ A_{11} & A_{12} & A_{13} \\ (A_{12})^T & A_{22} & A_{23} \\ (A_{13})^T & (A_{23})^T & A_{33} \end{pmatrix}.$$

By the form of the atoms, we know that $\widetilde{\mathcal{M}}(k; \beta^{(\ell)})$ and $\widetilde{\mathcal{M}}(k; \beta^{(c)})$ will be of the forms

$$(4.9) \quad \widetilde{\mathcal{M}}(k; \beta^{(c)}) = \begin{pmatrix} (\vec{X}^{(0,k)})^T \\ (\mathcal{T} \setminus \vec{X}^{(0,k)})^T \\ (\mathcal{C} \setminus (\vec{X}^{(0,k)} \cup \mathcal{T}))^T \end{pmatrix} \begin{pmatrix} \vec{X}^{(0,k)} & \mathcal{T} \setminus \vec{X}^{(0,k)} & \mathcal{C} \setminus (\vec{X}^{(0,k)} \cup \mathcal{T}) \\ A & A_{12} & A_{13} \\ (A_{12})^T & A_{22} & A_{23} \\ (A_{13})^T & (A_{23})^T & A_{33} \end{pmatrix},$$

$$\widetilde{\mathcal{M}}(k; \beta^{(\ell)}) = \begin{pmatrix} (\vec{X}^{(0,k)})^T \\ (\mathcal{T} \setminus \vec{X}^{(0,k)})^T \\ (\mathcal{C} \setminus (\vec{X}^{(0,k)} \cup \mathcal{T}))^T \end{pmatrix} \begin{pmatrix} \vec{X}^{(0,k)} & \mathcal{T} \setminus \vec{X}^{(0,k)} & \mathcal{C} \setminus (\vec{X}^{(0,k)} \cup \mathcal{T}) \\ A_{11} - A & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} \end{pmatrix}$$

for some Hankel matrix $A \in S_{k+1}$. Define matrix functions $\mathcal{F} : S_{k+1} \rightarrow S_{\frac{(k+1)(k+2)}{2}}$ and $\mathcal{H} : S_{k+1} \rightarrow S_{k+1}$ by

$$(4.10) \quad \mathcal{F}(\mathbf{A}) = \begin{pmatrix} \mathbf{A} & A_{12} & A_{13} \\ (A_{12})^T & A_{22} & A_{23} \\ (A_{13})^T & (A_{23})^T & A_{33} \end{pmatrix} \quad \text{and} \quad \mathcal{H}(\mathbf{A}) = A_{11} - \mathbf{A}.$$

Using (4.9), (4.7) becomes equivalent to

$$(4.11) \quad \widetilde{\mathcal{M}}(k; \beta) = \mathcal{F}(A) + \mathcal{H}(A) \oplus \mathbf{0}_{\frac{k(k+1)}{2}}$$

for some Hankel matrix $A \in S_{k+1}$.

Lemma 4.1. *Assume the notation above. The sequence $\beta = \{\beta_i\}_{i \in \mathbb{Z}_+^2, |i| \leq 2k}$, where $k \geq 3$, has a $\mathcal{Z}(p)$ -representing measure if and only if there exist a Hankel matrix $A \in S_{k+1}$, such that:*

- (1) *The sequence with the moment matrix $\mathcal{F}(A)$ has a $\mathcal{Z}(c)$ -representing measure.*
- (2) *The sequence with the moment matrix $\mathcal{H}(A)$ has a \mathbb{R} -representing measure.*

Proof. First we prove the implication (\Rightarrow). If β has a $\mathcal{Z}(p)$ -rm μ , then μ is supported on the union of the line $y = 0$ and the conic $c(x, y) = 0$. Since the moment matrix, generated by the measure supported on $y = 0$, can be nonzero only when restricted to the columns and rows indexed by $\vec{X}^{(0,k)}$, it follows that the moment matrix generated by the restriction $\mu|_{\{c=0\}}$ (resp. $\mu|_{\{y=0\}}$) of the measure μ to the conic $c(x, y) = 0$ (resp. line $y = 0$), is of the form $\mathcal{F}(A)$ (resp. $\mathcal{H}(A) \oplus \mathbf{0}_{\frac{k(k+1)}{2}}$) for some Hankel matrix $A \in S_{k+1}$.

It remains to establish the implication (\Leftarrow). Let $\mathcal{M}^{(c)}(k)$ (resp. $\mathcal{M}^{(\ell)}(k)$) be the moment matrix generated by the measure μ_1 (resp. μ_2) supported on $\mathcal{Z}(c)$ (resp. $y = 0$) such that

$$(4.12) \quad P\mathcal{M}^{(c)}(k)P^T = \mathcal{F}(A), \quad P\mathcal{M}^{(\ell)}(k)P^T = \mathcal{H}(A) \oplus \mathbf{0}_{\frac{k(k+1)}{2}},$$

respectively, where P is as in (4.6). The equalities (4.12) imply that $\mathcal{M}(k; \beta) = \mathcal{M}^{(c)}(k) + \mathcal{M}^{(\ell)}(k; \beta)$. Since the measure $\mu_1 + \mu_2$ is supported on the curve $\mathcal{Z}(c) \cup \{y = 0\} = \mathcal{Z}(p)$, the implication (\Leftarrow) holds. \square

Lemma 4.2. *Assume the notation above and let the sequence $\beta = \{\beta_i\}_{i \in \mathbb{Z}_+^2, |i| \leq 2k}$, where $k \geq 3$, admit a $\mathcal{Z}(p)$ -representing measure. Let $A := A_{(\beta_{0,0}^{(c)}, \beta_{1,0}^{(c)}, \dots, \beta_{2k,0}^{(c)})} \in S_{k+1}$ be a Hankel matrix such that $\mathcal{F}(A)$ admits a $\mathcal{Z}(c)$ -representing measure and $\mathcal{H}(A)$ admits a \mathbb{R} -representing measure. Let $c(x, y)$ be of the form*

$$(4.13) \quad c(x, y) = a_{00} + a_{10}x + a_{20}x^2 + a_{01}y + a_{02}y^2 + a_{11}xy \quad \text{with } a_{ij} \in \mathbb{R}$$

and exactly one of the coefficients a_{00}, a_{10}, a_{20} is nonzero.

If:

(1) $a_{00} \neq 0$, then

$$\beta_{i,0}^{(c)} = -\frac{1}{a_{00}}(a_{01}\beta_{i,1} + a_{02}\beta_{i,2} + a_{11}\beta_{i+1,1}) \quad \text{for } i = 0, \dots, 2k - 2.$$

(2) $a_{10} \neq 0$, then

$$\beta_{i,0}^{(c)} = -\frac{1}{a_{10}}(a_{01}\beta_{i,1} + a_{02}\beta_{i,2} + a_{11}\beta_{i+1,1}) \quad \text{for } i = 1, \dots, 2k - 1.$$

(3) $a_{20} \neq 0$, then

$$\beta_{i,0}^{(c)} = -\frac{1}{a_{20}}(a_{01}\beta_{i,1} + a_{02}\beta_{i,2} + a_{11}\beta_{i+1,1}) \quad \text{for } i = 2, \dots, 2k.$$

Proof. By Lemma 4.1, $\mathcal{F}(A)$ has a $\mathcal{Z}(c)$ -rm for some Hankel matrix $A \in S_{k+1}$. Hence, $\mathcal{F}(A)$ satisfies the rg relations $X^i Y^j c(X, Y) = \mathbf{0}$ for $i, j \in \mathbb{Z}_+$, $i + j \leq k - 2$. Let us assume that $a_{00} \neq 0$ and $a_{10} = a_{20} = 0$. In particular, $\mathcal{F}(A)$ satisfies the relations

$$(4.14) \quad \begin{aligned} a_{00}1 + a_{01}Y + a_{02}Y^2 + a_{11}XY &= \mathbf{0}, \\ a_{00}X^{k-2} + a_{01}X^{k-2}Y + a_{02}X^{k-2}Y^2 + a_{11}X^{k-1}Y &= \mathbf{0}. \end{aligned}$$

Observing the rows $1, X, \dots, X^k$ of $\mathcal{F}(A)$, the relations (4.14) imply that

$$(4.15) \quad \beta_{i,0}^{(c)} = -\frac{1}{a_{00}}(a_{01}\beta_{i,1}^{(c)} + a_{02}\beta_{i,2}^{(c)} + a_{11}\beta_{i+1,1}^{(c)}) \quad \text{for } i = 0, \dots, 2k - 2.$$

Using the forms of $\widetilde{\mathcal{M}}(k; \beta)$ and $\mathcal{F}(A)$ (see (4.8) and (4.10)), it follows that $\beta_{i,1}^{(c)} = \beta_{i,1}$ and $\beta_{j,2}^{(c)} = \beta_{j,2}$ for each i, j . Using this in (4.15) proves the statement (1) of the lemma. The proofs of the statements (2) and (3) are analogous. \square

Lemma 4.2 states that for all canonical relations from Proposition 3.1 except for the mixed type relation, all but two entries of the Hankel matrix A from Lemma 4.1 are uniquely determined by β . The following lemma gives the smallest candidate for A in Lemma 4.1 with respect to the usual Loewner order of matrices.

Lemma 4.3. *Assume the notation above and let $\beta = \{\beta_i\}_{i \in \mathbb{Z}_+^2, |i| \leq 2k}$, where $k \geq 3$, be a sequence of degree $2k$. Assume that $\widetilde{\mathcal{M}}(k; \beta)$ is positive semidefinite and satisfies the column relations (4.5). Then:*

(1) $\mathcal{F}(A) \succeq 0$ for some $A \in S_{k+1}$ if and only if $A \succeq A_{12}(A_{22})^\dagger(A_{12})^T$.

(2) $\mathcal{F}(A_{12}(A_{22})^\dagger(A_{12})^T) \succeq 0$ and $\mathcal{H}(A_{12}(A_{22})^\dagger(A_{12})^T) \succeq 0$.

(3) $\mathcal{F}(A_{12}(A_{22})^\dagger(A_{12})^T)$ satisfies the column relations $X^i Y^j c(X, Y) = 0$ for $i, j \in \mathbb{Z}_+$ such that $i + j \leq k - 2$.

(4) We have that

$$\begin{aligned} \text{rank } \widetilde{\mathcal{M}}(k; \beta) &= \text{rank } A_{22} + \text{rank } (A_{11} - A_{12}(A_{22})^\dagger(A_{12})^T) \\ &= \text{rank } \mathcal{F}(A_{12}(A_{22})^\dagger(A_{12})^T) + \text{rank } \mathcal{H}(A_{12}(A_{22})^\dagger(A_{12})^T). \end{aligned}$$

Proof. By the equivalence between (1a) and (1b) of Theorem 2.2 used for $(M, A) = (\widetilde{\mathcal{M}}(k; \beta), A_{11})$ and $(M, A) = ((\widetilde{\mathcal{M}}(k; \beta))_{\vec{X}(0,k) \cup \mathcal{T}}, A_{11})$, it follows in particular that

$$(4.16) \quad \begin{aligned} \mathcal{C}\left(\begin{pmatrix} (A_{12})^T \\ (A_{13})^T \end{pmatrix}\right) &\subseteq \mathcal{C}\left(\begin{pmatrix} A_{22} & A_{23} \\ (A_{23})^T & A_{33} \end{pmatrix}\right), \\ \mathcal{C}(A_{12}^T) &\subseteq \mathcal{C}(A_{22}). \end{aligned}$$

and

$$(4.17) \quad \mathcal{H}(A_{\min}) \succeq 0,$$

where

$$A_{\min} := (A_{12} \quad A_{13}) \begin{pmatrix} A_{22} & A_{23} \\ (A_{23})^T & A_{33} \end{pmatrix}^\dagger \begin{pmatrix} (A_{12})^T \\ (A_{13})^T \end{pmatrix}.$$

Using the equivalence between (1a) and (1b) of Theorem 2.2 again for the pairs $(M, A) = (\mathcal{F}(A), A)$ and $(M, A) = ((\mathcal{F}(A))_{\vec{X}(0,k) \cup \mathcal{T}}, A)$, it follows that

$$(4.18) \quad \begin{aligned} \mathcal{F}(A) \succeq 0 &\Leftrightarrow A \succeq A_{\min}, \\ (\mathcal{F}(A))_{\vec{X}(0,k) \cup \mathcal{T}} \succeq 0 &\Leftrightarrow A \succeq A_{12}(A_{22})^\dagger(A_{12})^T =: \widetilde{A}_{\min}. \end{aligned}$$

Since $\mathcal{F}(A) \succeq 0$ implies, in particular, that $(\mathcal{F}(A))_{\vec{X}(0,k) \cup \mathcal{T}} \succeq 0$, (4.18) implies that

$$(4.19) \quad A_{\min} \succeq \widetilde{A}_{\min}.$$

Claim. $A_{\min} = \widetilde{A}_{\min}$.

Proof of Claim. By (4.18) and (4.19), it suffices to prove that $\mathcal{F}(\widetilde{A}_{\min}) \succeq 0$. By definition of \mathcal{T} and the relations $X^i Y^j p(X, Y) = X^i Y^{j+1} c(X, Y) = \mathbf{0}$, $i, j \in \mathbb{Z}_+$, $i + j \leq k - 3$, which hold in $\widetilde{\mathcal{M}}(k; \beta)$, it follows, in particular, that

$$(4.20) \quad \mathcal{C}\left(\begin{pmatrix} A_{23} \\ A_{33} \end{pmatrix}\right) \subseteq \mathcal{C}\left(\begin{pmatrix} A_{22} \\ (A_{23})^T \end{pmatrix}\right)$$

(4.16) and (4.20) together imply that

$$(4.21) \quad \mathcal{C}\left(\begin{pmatrix} (A_{12})^T \\ (A_{13})^T \end{pmatrix}\right) \subseteq \mathcal{C}\left(\begin{pmatrix} A_{22} \\ (A_{23})^T \end{pmatrix}\right).$$

(4.16) and (4.21) can be equivalently expressed as

$$(4.22) \quad \begin{aligned} \begin{pmatrix} A_{22} \\ (A_{23})^T \end{pmatrix} W &= \begin{pmatrix} A_{23} \\ A_{33} \end{pmatrix} \text{ for some matrix } W, \\ \begin{pmatrix} A_{22} \\ (A_{23})^T \end{pmatrix} X &= \begin{pmatrix} (A_{12})^T \\ (A_{13})^T \end{pmatrix} \text{ for some matrix } X. \end{aligned}$$

We have that

$$\begin{aligned} 0 &\preceq \begin{pmatrix} X^T \\ I \\ W^T \end{pmatrix} A_{22} \begin{pmatrix} X & I & W \end{pmatrix} \\ &= \begin{pmatrix} X^T A_{22} X & X^T A_{22} & X^T A_{22} W \\ A_{22} X & A_{22} & A_{22} W \\ W^T A_{22} X & W^T A_{22} & W^T A_{22} W \end{pmatrix} \\ &= \begin{pmatrix} A_{12}(A_{22})^\dagger(A_{12})^T & A_{12} & A_{13} \\ (A_{12})^T & A_{22} & A_{23} \\ (A_{13})^T & (A_{23})^T & A_{33} \end{pmatrix} = \mathcal{F}(\tilde{A}_{\min}) \end{aligned}$$

where I is the identity matrix of the same size as A_{22} and we used (4.22) in the second equality. This proves the Claim. \blacksquare

Using (4.17), (4.18) and Claim, the statements (1) and (2) follow. By Theorem 2.2.(2), used for $(M, A) = (\tilde{\mathcal{M}}(k; \beta), A_{11})$, we have that

$$(4.23) \quad \begin{aligned} \text{rank } \tilde{\mathcal{M}}(k; \beta) &= \text{rank} \begin{pmatrix} A_{22} & A_{23} \\ (A_{23})^T & A_{33} \end{pmatrix} + \text{rank } \mathcal{H}(A_{\min}) \\ &= \text{rank } \mathcal{F}(A_{\min}) + \text{rank } \mathcal{H}(A_{\min}). \end{aligned}$$

By (4.20) and

$$B := \begin{pmatrix} A_{22} & A_{23} \\ (A_{23})^T & A_{33} \end{pmatrix} \succeq 0,$$

it follows by Theorem 2.2, used for $(M, A) = (B, A_{22})$, that $\text{rank } B = \text{rank } A_{22}$. Using this and the Claim, (4.23) implies the statement (4).

Since $\tilde{\mathcal{M}}(k; \beta)$ satisfies the relations (4.5), it follows that the restriction $(\mathcal{F}(\tilde{A}_{\min}))_{\mathcal{C} \setminus \vec{X}^{(0,k)}, \mathcal{C}}$ satisfies the column relations $X^i Y^j \mathcal{C}(X, Y) = \mathbf{0}$ for $i, j \in \mathbb{Z}_+$ such that $i + j \leq k - 2$. By Proposition 2.3, these relations extend to $\mathcal{F}(\tilde{A}_{\min})$, which proves (3). \square

Remark 4.4. By Lemmas 4.1–4.3, solving the $\mathcal{Z}(p)$ –TMP for the sequence $\beta = \{\beta_i\}_{i \in \mathbb{Z}_+, |i| \leq 2k}$, where $k \geq 3$, with p being any but the mixed type relation from Proposition 3.1, the natural procedure is the following:

- (1) First compute $A_{\min} := A_{12}(A_{22})^\dagger A_{12}$. By Lemma 4.3.(3), there is one entry of A_{\min} , which might need to be changed to obtain a Hankel structure. Namely, in the notation (4.13), if:
 - (a) $a_{00} \neq 0$, then the value of $(A_{\min})_{k,k}$ must be made equal to $(A_{\min})_{k-1,k+1}$.
 - (b) $a_{10} \neq 0$, then the value of $(A_{\min})_{1,k+1}$ must be made equal to $(A_{\min})_{2,k}$.
 - (c) $a_{20} \neq 0$, then the value of $(A_{\min})_{2,2}$ must be made equal to $(A_{\min})_{3,1}$.

Let \widehat{A}_{\min} be the matrix obtained from A_{\min} after performing the changes described above.

- (2) Study if $\mathcal{F}(\widehat{A}_{\min})$ and $\mathcal{H}(\widehat{A}_{\min})$ admit a $\mathcal{Z}(c)$ -rm and a \mathbb{R} -rm, respectively. If the answer is yes, β admits a $\mathcal{Z}(p)$ -rm. Otherwise by Lemma 4.2, there are two antidiagonals of the Hankel matrix \widehat{A}_{\min} , which can be varied so that the matrices $\mathcal{F}(\widehat{A}_{\min})$ and $\mathcal{H}(\widehat{A}_{\min})$ will admit the corresponding measures. Namely, in the notation (4.13), if:

- (a) $a_{00} \neq 0$, then the last two antidiagonals of \widehat{A}_{\min} can be changed.
- (b) $a_{10} \neq 0$, then the left-upper and the right-lower corner of \widehat{A}_{\min} can be changed.
- (c) $a_{20} \neq 0$, then the first two antidiagonals of \widehat{A}_{\min} can be changed.

To solve the $\mathcal{Z}(p)$ -TMP for β one needs to characterize, when it is possible to change these antidiagonals in such a way to obtain a matrix \check{A}_{\min} , such that $\mathcal{F}(\check{A}_{\min})$ and $\mathcal{H}(\check{A}_{\min})$ admit a $\mathcal{Z}(c)$ -rm and a \mathbb{R} -rm, respectively.

In Sections 5 and 6 we solve concretely the TMP on reducible cubic curves in the circular and parabolic type form (see the classification from Proposition 3.1). The parallel lines type form was solved in [Zal22a], while the hyperbolic type forms will be solved in the forthcoming work [YZ+].

5. CIRCULAR TYPE RELATION: $p(x, y) = y(ay + x^2 + y^2)$, $a \notin \mathbb{R} \setminus \{0\}$.

In this section we solve the $\mathcal{Z}(p)$ -TMP for the sequence $\beta = \{\beta_{i,j}\}_{i,j \in \mathbb{Z}_+, i+j \leq 2k}$ of degree $2k$, $k \geq 3$, where $p(x, y) = y(ay + x^2 + y^2)$, $a \in \mathbb{R} \setminus \{0\}$. Assume the notation from Section 4. If β admits a $\mathcal{Z}(p)$ -TMP, then $\mathcal{M}(k; \beta)$ must satisfy the relations

$$(5.1) \quad aY^{2+j}X^i + Y^{1+j}X^{2+i} = -Y^{3+j}X^i \quad \text{for } i, j \in \mathbb{Z}_+ \text{ such that } i + j \leq k - 3.$$

In the presence of all column relations (5.1), the column space $\mathcal{C}(\mathcal{M}(k; \beta))$ is spanned by the columns in the set

$$(5.2) \quad \mathcal{T} = \vec{X}^{(0,k)} \cup Y\vec{X}^{(0,k-1)} \cup Y^2\vec{X}^{(0,k-2)},$$

where

$$Y^i\vec{X}^{(j,\ell)} := (Y^iX^j, Y^iX^{j+1}, \dots, Y^iX^\ell) \quad \text{with } i, j, \ell \in \mathbb{Z}_+, j \leq \ell, i + \ell \leq k.$$

Let $\widetilde{\mathcal{M}}(k; \beta)$ be as in (4.9). Let

$$(5.3) \quad A_{\min} := A_{12}(A_{22})^\dagger(A_{12})^T.$$

As described in Remark 4.4, A_{\min} might need to be changed to

$$\widehat{A}_{\min} = A_{\min} + \eta E_{2,2}^{(k+1)},$$

where

$$\eta := (A_{\min})_{1,3} - (A_{\min})_{2,2}.$$

Let $\mathcal{F}(\mathbf{A})$ and $\mathcal{H}(\mathbf{A})$ be as in (4.10). Write

$$\mathcal{H}(\widehat{A}_{\min}) := \begin{matrix} & & 1 & & X & & \vec{X}^{(2,k)} \\ & & & & & & \\ & 1 & & & & & \\ & X & & & & & \\ (\vec{X}^{(2,k)})^T & & \begin{pmatrix} \beta_{0,0} - (A_{\min})_{1,1} & \beta_{1,0} - (A_{\min})_{1,2} & (h_{12}^{(1)})^T \\ \beta_{1,0} - (A_{\min})_{1,2} & \beta_{2,0} - (A_{\min})_{1,3} & (h_{12}^{(2)})^T \\ h_{12}^{(1)} & h_{12}^{(2)} & H_{22} \end{pmatrix} & & & & \end{matrix},$$

$$(5.4) \quad H_1 := (\mathcal{H}(\widehat{A}_{\min}))_{\{1\} \cup \vec{X}^{(2,k)}} = \begin{matrix} & & 1 & & \vec{X}^{(2,k)} \\ & & & & \\ & 1 & & & \\ & (\vec{X}^{(2,k)})^T & & & \\ & & \begin{pmatrix} \beta_{0,0} - (A_{\min})_{1,1} & (h_{12}^{(1)})^T \\ h_{12}^{(1)} & H_{22} \end{pmatrix} & & \end{matrix},$$

$$H_2 := (\mathcal{H}(\widehat{A}_{\min}))_{\vec{X}^{(1,k)}} = \begin{matrix} & & X & & \vec{X}^{(2,k)} \\ & & & & \\ & X & & & \\ (\vec{X}^{(2,k)})^T & & \begin{pmatrix} \beta_{2,0} - (A_{\min})_{1,3} & (h_{12}^{(2)})^T \\ h_{12}^{(2)} & H_{22} \end{pmatrix} & & \end{matrix}.$$

Define also the matrix function

$$(5.5) \quad \mathcal{G} : \mathbb{R}^2 \rightarrow S_{k+1}, \quad \mathcal{G}(\mathbf{t}, \mathbf{u}) = \widehat{A}_{\min} + \mathbf{t}E_{1,1}^{(k+1)} + \mathbf{u}(E_{1,2}^{(k+1)} + E_{2,1}^{(k+1)}).$$

The solution to the cubic circular type relation TMP is the following.

Theorem 5.1. *Let $p(x, y) = y(ay + x^2 + y^2)$, $a \in \mathbb{R} \setminus \{0\}$, and $\beta = (\beta_{i,j})_{i,j \in \mathbb{Z}_+, i+j \leq 2k}$, where $k \geq 3$. Assume also the notation above. Then the following statements are equivalent:*

- (1) β has a $\mathcal{Z}(p)$ -representing measure.
- (2) $\widetilde{\mathcal{M}}(k; \beta)$ is positive semidefinite, the relations

$$(5.6) \quad a\beta_{i,2+j} + \beta_{2+i,1+j} = -\beta_{i,3+j} \quad \text{hold for every } i, j \in \mathbb{Z}_+ \text{ with } i + j \leq 2k - 3$$

and one of the following statements holds:

- (a) $\eta = 0$ and one of the following holds:

- (i) $\text{rank}(\mathcal{H}(A_{\min}))_{\vec{X}^{(0,k-1)}} = k$.
- (ii) $\text{rank}(H_2)_{\vec{X}^{(1,k-1)}} = \text{rank } H_2$.

- (b) $\eta > 0$, H_2 is positive semidefinite and defining a real number

$$(5.7) \quad u_0 = \beta_{1,0} - (A_{\min})_{1,2} - (h_{12}^{(1)})^T (H_{22})^\dagger h_{12}^{(2)},$$

a function

$$(5.8) \quad h(\mathbf{t}) = \sqrt{(H_1/H_{22} - \mathbf{t})(H_2/H_{22})}$$

and a set

$$(5.9) \quad \mathcal{I} = \left\{ (t, \sqrt{\eta t}) \in \mathbb{R}_+ \times \mathbb{R}_+ : \sqrt{\eta t} = u_0 + h(t) \right\}, \\ \cup \left\{ (t, \sqrt{\eta t}) \in \mathbb{R}_+ \times \mathbb{R}_- : \sqrt{\eta t} = u_0 - h(t) \right\}, \\ \cup \left\{ (t, -\sqrt{\eta t}) \in \mathbb{R}_+ \times \mathbb{R}_+ : -\sqrt{\eta t} = u_0 + h(t) \right\}, \\ \cup \left\{ (t, -\sqrt{\eta t}) \in \mathbb{R}_+ \times \mathbb{R}_- : -\sqrt{\eta t} = u_0 - h(t) \right\},$$

one of the following holds:

- (i) The set \mathcal{I} has two elements and H_2 is positive definite.

(ii) $\mathcal{I} = \{(\tilde{t}, \tilde{u})\}$ and

$$(5.10) \quad \text{rank} \left((\mathcal{H}(\mathcal{G}(\tilde{t}, \tilde{u})))_{\tilde{\mathcal{X}}(0, k-1)} \right) = \text{rank} \mathcal{H}(\mathcal{G}(\tilde{t}, \tilde{u})).$$

Moreover, if a $\mathcal{Z}(p)$ -representing measure for β exists, then:

- There exists at most $(\text{rank } \widetilde{\mathcal{M}}(k; \beta) + 1)$ -atomic $\mathcal{Z}(p)$ -representing measure.
- There exists a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta))$ -atomic $\mathcal{Z}(p)$ -representing measure if and only if any of the following holds:
 - $\eta = 0$.
 - $\eta > 0$ and $\mathcal{H}(A_{\min})$ is positive definite.

In particular, a p -pure sequence β with a $\mathcal{Z}(p)$ -representing measure admits a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta))$ -atomic $\mathcal{Z}(p)$ -representing measure.

Remark 5.2. In this remark we explain the idea of the proof of Theorem 5.1 and the meaning of the conditions in the statement of the theorem.

By Lemmas 4.1–4.2, the existence of a $\mathcal{Z}(p)$ -rm for β is equivalent to the existence of $t, u \in \mathbb{R}$ such that $\mathcal{F}(\mathcal{G}(t, u))$ admits a $\mathcal{Z}(ay + x^2 + y^2)$ -rm and $\mathcal{H}(\mathcal{G}(t, u))$ admits a \mathbb{R} -rm. Let

$$\mathcal{R}_1 = \{(t, u) \in \mathbb{R}^2 : \mathcal{F}(\mathcal{G}(t, u)) \succeq 0\} \quad \text{and} \quad \mathcal{R}_2 = \{(t, u) \in \mathbb{R}^2 : \mathcal{H}(\mathcal{G}(t, u)) \succeq 0\}.$$


We denote by ∂R_i and \mathring{R}_i the topological boundary and the interior of the set R_i , respectively. By the necessary conditions for the existence of a $\mathcal{Z}(p)$ -rm [CF04, Fia95, CF96], $\widetilde{\mathcal{M}}(k; \beta)$ must be psd and the relations (5.6) must hold. Using also Theorem 2.6, Theorem 5.1.(1) is equivalent to

$$(5.11) \quad \begin{aligned} &\widetilde{\mathcal{M}}(k; \beta) \succeq 0, \text{ the relations (5.6) hold and} \\ &\exists (t_0, u_0) \in \mathcal{R}_1 \cap \mathcal{R}_2 : \mathcal{H}(\mathcal{G}(t_0, u_0)) \text{ admits a } \mathbb{R}\text{-rm.} \end{aligned}$$

In the proof of Theorem 5.1 we show that (5.11) is equivalent to Theorem 5.1.(2):


(1) First we establish (see Claims 1 and 2 below) that the form of:

- \mathcal{R}_1 is one of the following:



where the left case occurs if $\eta > 0$ and the right if $\eta = 0$. The case $\eta < 0$ cannot occur.

- \mathcal{R}_2 is one of the following:



where the left case occurs if $H_2/H_{22} > 0$ and the right if $H_2/H_{22} = 0$.

(2) If $\eta = 0$, then we show that (5.11) is equivalent to

$\widetilde{\mathcal{M}}(k; \beta) \succeq 0$, the relations (5.6) hold and $\mathcal{H}(\mathcal{G}(0, 0))$ admits a \mathbb{R} -rm.

The latter statement is further equivalent to Theorem 5.1.(2a).

(3) If $\eta > 0$, then by the forms of \mathcal{R}_1 and \mathcal{R}_2 , $\mathcal{I} = \partial\mathcal{R}_1 \cap \partial\mathcal{R}_2$ is one of the following: (i) \emptyset , (ii) a one-element set, (iii) a two-element set. In the case:

- (i), a $\mathcal{Z}(p)$ -rm for β clearly cannot exist.
- (ii), then denoting $\mathcal{I} = \{(\tilde{t}, \tilde{u})\}$, (5.11) is equivalent to

$\widetilde{\mathcal{M}}(k; \beta) \succeq 0$, the relations (5.6) hold and $\mathcal{H}(\mathcal{G}(\tilde{t}, \tilde{u}))$ admits a \mathbb{R} -rm.

The latter statement is equivalent to Theorem 5.1.(2(b)ii).

- (iii), (5.11) is equivalent to H_2 being positive definite, which is Theorem 5.1.(2(b)i). Moreover, in this case for at least one of the points $(t, u) \in \mathcal{I}$, a $\mathcal{Z}(ay + x^2 + y^2)$ -rm and a \mathbb{R} -rm exist for $\mathcal{F}(\mathcal{G}(t, u))$ and $\mathcal{H}(\mathcal{G}(t, u))$, respectively.

Proof of Theorem 5.1. Let $\mathcal{R}_1, \mathcal{R}_2$ be as in Remark 5.2. As explained in Remark 5.2, Theorem 5.1.(1) is equivalent to (5.11), thus it remains to prove that (5.11) is equivalent to Theorem 5.1.(2).

First we establish a few claims needed in the proof. Claim 1 (resp. 2) describes \mathcal{R}_1 (resp. \mathcal{R}_2) concretely.

Claim 1. Assume that $\widetilde{\mathcal{M}}(k; \beta) \succeq 0$. Then

$$(5.12) \quad \mathcal{R}_1 = \begin{cases} \{(t, u) \in \mathbb{R}^2 : t \geq 0, u \in [-\sqrt{\eta t}, \sqrt{\eta t}]\}, & \text{if } \eta \geq 0, \\ \emptyset, & \text{if } \eta < 0. \end{cases}$$

If $\eta \geq 0$, we have

$$(5.13) \quad \text{rank } \mathcal{F}(\mathcal{G}(t, u)) = \begin{cases} \text{rank } \mathcal{F}(A_{\min}), & \text{if } t = 0, \eta = 0, \\ \text{rank } \mathcal{F}(A_{\min}) + 1, & \text{if } (t > 0 \text{ or } \eta > 0) \text{ and } u \in \{-\sqrt{\eta t}, \sqrt{\eta t}\}, \\ \text{rank } \mathcal{F}(A_{\min}) + 2, & \text{if } t > 0, \eta > 0, u \in (-\sqrt{\eta t}, \sqrt{\eta t}), \end{cases}$$

where A_{\min} is an in (5.3).

Proof of Claim 1. Note that

$$(5.14) \quad \begin{aligned} \mathcal{G}(\mathbf{t}, \mathbf{u}) &= A_{\min} + \eta E_{2,2}^{(k+1)} + \mathbf{t} E_{1,1}^{(k+1)} + \mathbf{u} (E_{1,2}^{(k+1)} + E_{2,1}^{(k+1)}) \\ &= A_{\min} + \begin{pmatrix} \mathbf{t} & \mathbf{u} \\ \mathbf{u} & \eta \end{pmatrix} \oplus \mathbf{0}_{k-1}. \end{aligned}$$

By Lemma 4.3, we have that

$$(5.15) \quad \mathcal{F}(\mathcal{G}(t, u)) \succeq 0 \Leftrightarrow \mathcal{G}(t, u) \succeq A_{\min}$$

Using (5.14), (5.15) and the definition of \mathcal{R}_1 , we have that

$$(5.16) \quad (t, u) \in \mathcal{R}_1 \Leftrightarrow \begin{pmatrix} t & u \\ u & \eta \end{pmatrix} \succeq 0 \Leftrightarrow t \geq 0, \eta \geq 0, t\eta \geq u^2,$$

which proves (5.12).

To prove (5.13) first note that by construction of $\mathcal{F}(A_{\min})$, the columns 1 and X are in the span of the columns indexed by $\mathcal{C} \setminus \vec{X}^{(0,k)}$. Hence, there are vectors

$$(5.17) \quad v_1, v_2 \in \ker \mathcal{F}(A_{\min})$$

of the forms

$$v_1 = (1 \quad \mathbf{0}_{1,k} \quad (\tilde{v}_1)^T)^T \in \mathbb{R}^{\frac{(k+1)(k+2)}{2}} \quad \text{and} \quad v_2 = (0 \quad 1 \quad \mathbf{0}_{1,k-2} \quad (\tilde{v}_2)^T)^T \in \mathbb{R}^{\frac{(k+1)(k+2)}{2}}.$$

Let $r := \text{rank} \begin{pmatrix} t & u \\ u & \eta \end{pmatrix}$. Clearly,

$$(5.18) \quad \text{rank} \mathcal{F}(\mathcal{G}(t, u)) \leq \text{rank} \mathcal{F}(A_{\min}) + r.$$

We separate three cases according to r .

Case 1: $r = 0$. In this case $t = u = \eta = 0$ and $\mathcal{G}(0, 0) = A_{\min}$. In this case (5.13) clearly holds.

Case 2: $r = 1$. In this case $t\eta = u^2$. Together with (5.16), this is equivalent to $(t > 0$ or $\eta > 0)$ and $u \in \{-\sqrt{\eta t}, \sqrt{\eta t}\}$. By (5.18) and $\mathcal{F}(\mathcal{G}(t, u)) \succeq \mathcal{F}(A_{\min})$ to prove (5.13), it suffices to find $v \in \ker \mathcal{F}(A_{\min})$ and $v \notin \ker \mathcal{F}(\mathcal{G}(t, u))$. Note that at least one of v_1, v_2 from (5.17) is such a vector, since

$$(v_1)^T \mathcal{F}(\mathcal{G}(t, u)) v_1 = t \quad \text{and} \quad (v_2)^T \mathcal{F}(\mathcal{G}(t, u)) v_2 = \eta.$$

Case 3: $r = 2$. In this case $t\eta > u^2$. Together with (5.16), this is equivalent to $t > 0, \eta > 0, u \in (-\sqrt{\eta t}, \sqrt{\eta t})$. Note that

$$(5.19) \quad \mathcal{F}(\mathcal{G}(t, u)) = \mathcal{F}\left(\mathcal{G}\left(\frac{u^2}{\eta}, u\right)\right) + \left(t - \frac{u^2}{\eta}\right) \oplus \mathbf{0}_{\frac{(k+1)(k+2)}{2}-1} \succeq \mathcal{F}\left(\mathcal{G}\left(\frac{u^2}{\eta}, u\right)\right).$$

By Case 2, we have $\text{rank} \mathcal{F}\left(\mathcal{G}\left(\frac{u^2}{\eta}, u\right)\right) = \text{rank} \mathcal{F}(A_{\min}) + 1$. By (5.18) and (5.19), to prove (5.13), it suffices to find $v \in \ker \mathcal{F}\left(\mathcal{G}\left(\frac{u^2}{\eta}, u\right)\right)$ and $v \notin \ker \mathcal{F}(\mathcal{G}(t, u))$. We will check below, that v_3 , defined by

$$v_3 = v_1 - \frac{u}{\eta} v_2 = \left(1 \quad -\frac{u}{\eta} \quad (\tilde{v}_3)^T\right)^T \in \mathbb{R}^{\frac{(k+1)(k+2)}{2}},$$

is such a vector. This follows by

$$\mathcal{F}\left(\mathcal{G}\left(\frac{u^2}{\eta}, u\right)\right) v_3 = \mathcal{F}(A_{\min}) v_3 + \left(\begin{pmatrix} \frac{u^2}{\eta} & u \\ u & \eta \end{pmatrix} \oplus \mathbf{0}_{\frac{(k+1)(k+2)}{2}-1}\right) v_3 = \mathbf{0}_{\frac{(k+1)(k+2)}{2}, 1}$$

Proof of Claim 4. Note that the set \mathcal{I} is equal to $\partial\mathcal{R}_1 \cap \partial\mathcal{R}_2$ (see (5.12) and (5.20)). Further on, $\partial\mathcal{R}_1$ is the union of the square root functions $\pm\sqrt{\eta\mathbf{t}}$, defined for $\mathbf{t} \in [0, \infty)$. Similarly, $\partial\mathcal{R}_2$ is the union of the square root functions $u_0 \pm \sqrt{(H_1/H_{22} - \mathbf{t})(H_2/H_{22})}$, defined for $\mathbf{t} \in (-\infty, t_0]$. If $H_2/H_{22} = 0$, then the latter could be a half-line $\{(t, u_0) : t \leq t_0\}$. If $\mathcal{R}_1 \cap \mathcal{R}_2 \neq \emptyset$, then geometrically it is clear that \mathcal{I} contains one or two elements. Assume that \mathcal{I} contains only one element, denoted by (\tilde{t}, \tilde{u}) . Clearly, $\mathcal{I} \subseteq \mathcal{R}_1 \cap \mathcal{R}_2$. Further on, we either have $\mathcal{I} = \mathcal{R}_1 \cap \mathcal{R}_2$ or $\mathcal{I} \subsetneq \mathcal{R}_1 \cap \mathcal{R}_2$. By the forms of $\partial\mathcal{R}_1$ and $\partial\mathcal{R}_2$, the latter case occurs if $H_2/H_{22} = 0$ or equivalently $\partial\mathcal{R}_2 = \mathcal{R}_2 = \{(t, u_0) : t \leq t_0\}$. But then the whole line segment $\{(t, u_0) : \tilde{t} \leq t \leq t_0\}$ lies in \mathcal{R}_1 , which proves Claim 4. \blacksquare

Claim 5. Let H_2 (see (5.4)) be positive definite, $(t_1, u_1) \in \partial\mathcal{R}_2$, $(t_2, u_2) \in \partial\mathcal{R}_2$ and $u_1 \neq u_2$. Then at least one of $\mathcal{H}(\mathcal{G}(t_1, u_1))$ and $\mathcal{H}(\mathcal{G}(t_2, u_2))$ admits a \mathbb{R} -rm.

Proof of Claim 5. Note that $\mathcal{H}(\mathcal{G}(t_i, u_i))$, $i = 1, 2$, is of the form

$$\mathcal{H}(\mathcal{G}(t_i, u_i)) = \begin{array}{c} 1 \\ X \\ (\vec{X}^{(2,k-1)})^T \\ X^k \end{array} \begin{pmatrix} 1 & X & \vec{X}^{(2,k-1)} & X^k \\ \beta_{0,0} - (A_{\min})_{1,1} - t_i & \beta_{1,0} - (A_{\min})_{1,2} - u_i & (\hat{h}_{12}^{(1)})^T & \tilde{\beta}_{k,0} \\ \beta_{1,0} - (A_{\min})_{1,2} - u_i & \beta_{2,0} - (A_{\min})_{1,3} & (\hat{h}_{12}^{(2)})^T & \tilde{\beta}_{k+1,0} \\ \hat{h}_{12}^{(1)} & \hat{h}_{12}^{(2)} & \hat{H}_2 & \hat{h}_3 \\ \tilde{\beta}_{k,0} & \tilde{\beta}_{k+1,0} & (\hat{h}_3)^T & \tilde{\beta}_{2k,0} \end{pmatrix}.$$

Assume on the contrary that none of $\mathcal{H}(\mathcal{G}(t_1, u_1))$ and $\mathcal{H}(\mathcal{G}(t_2, u_2))$ admits a \mathbb{R} -rm. Theorem 2.5 implies that the column X^k of $\mathcal{H}(\mathcal{G}(t_i, u_i))$, $i = 1, 2$, is not in the span of the other columns. Using this fact, the facts that $\mathcal{H}(\mathcal{G}(t_i, u_i))$, $i = 1, 2$, are not pd (by $(t_i, u_i) \in \partial\mathcal{R}_2$, $i = 1, 2$) and H_2 is pd, it follows that there is a column relation $1 = \sum_{j=1}^{k-1} \alpha_j^{(i)} X^j$, $\alpha_j^{(i)} \in \mathbb{R}$, in $\mathcal{H}(\mathcal{G}(t_i, u_i))$, $i = 1, 2$. Since $\mathcal{H}(\mathcal{G}(t_i, u_i)) \succeq 0$, $i = 1, 2$, it follows in particular by Theorem 2.2, used for $(M, A) = (\mathcal{H}(\mathcal{G}(t_i, u_i)), (\mathcal{H}(\mathcal{G}(t_i, u_i)))_{\vec{X}^{(0,k-1)}})$, $i = 1, 2$, that

$$(5.22) \quad \begin{pmatrix} \tilde{\beta}_{k,0} & \tilde{\beta}_{k+1,0} & (\hat{h}_3)^T \end{pmatrix}^T \in \mathcal{C}\left(\left(\mathcal{H}(\mathcal{G}(t_i, u_i))\right)_{\vec{X}^{(0,k-1)}}\right), \quad i = 1, 2.$$

Since the first column of $\mathcal{H}(\mathcal{G}(t_i, u_i)) \succeq 0$, $i = 1, 2$, is in the span of the others, (5.22) is equivalent to

$$(5.23) \quad \begin{pmatrix} \tilde{\beta}_{k,0} & \tilde{\beta}_{k+1,0} & (\hat{h}_3)^T \end{pmatrix}^T \in \mathcal{C}\left(\left(\mathcal{H}(\mathcal{G}(t_i, u_i))\right)_{\vec{X}^{(0,k-1)}, \vec{X}^{(1,k-1)}}\right), \quad i = 1, 2.$$

Since

$$\tilde{H}_2 := \left(\mathcal{H}(\mathcal{G}(t_i, u_i))\right)_{\vec{X}^{(1,k-1)}}, \quad i = 1, 2,$$

is invertible as a principal submatrix of H_2 , it follows that

$$(5.24) \quad \begin{pmatrix} \tilde{\beta}_{k,0} & \tilde{\beta}_{k+1,0} & (\hat{h}_3)^T \end{pmatrix}^T = \left(\left(\mathcal{H}(\mathcal{G}(t_i, u_i))\right)_{\vec{X}^{(0,k-1)}, \vec{X}^{(1,k-1)}}\right) v, \quad i = 1, 2.$$

with

$$v = \tilde{H}_2^{-1} \begin{pmatrix} \tilde{\beta}_{k+1,0} & \hat{h}_3 \end{pmatrix}^T = (v_1 \ v_2 \ \cdots \ v_{k-1})^T.$$

If $v_1 \neq 0$, this contradicts to (5.24) since $u_1 \neq u_2$. Hence, $v_1 = 0$. By the Hankel structure of $\mathcal{H}(\mathcal{G}(t_i, u_i))$, $i = 1, 2$, we have that

$$\left(\mathcal{H}(\mathcal{G}(t_i, u_i))\right)_{\vec{X}^{(0,k-2)}, \vec{X}^{(2,k)}} = \left(\mathcal{H}(\mathcal{G}(t_i, u_i))\right)_{\vec{X}^{(1,k-1)}, \vec{X}^{(1,k-1)}}, \quad i = 1, 2.$$

Then (5.24) and $v_1 = 0$ imply that

$$(5.25) \quad \left((\mathcal{H}(\mathcal{G}(t_i, u_i)))_{\vec{X}(0, k-2), \vec{X}(2, k)} \right) \tilde{v} = \left((\mathcal{H}(\mathcal{G}(t_i, u_i)))_{\vec{X}(1, k-1), \vec{X}(1, k-1)} \right) \tilde{v} = \mathbf{0}_{k+1, 1},$$

where $\tilde{v} = (v_2 \ \cdots \ v_{k-1} \ -1)$. Since $(\mathcal{H}(\mathcal{G}(t_i, u_i)))_{\vec{X}(1, k-1), \vec{X}(1, k-1)}$, $i = 1, 2$, is a principal submatrix of H_2 , (5.25) contradicts to H_2 being pd. This proves Claim 5. \blacksquare

Now we prove the implication (5.11) \Rightarrow Theorem 5.1.(2). Since $(t_0, u_0) \in \mathcal{R}_1$, it follows that $\mathcal{R}_1 \neq \emptyset$. By (5.12), $\eta \geq 0$. We separate two cases according to the value of η .

Case 1: $\eta = 0$. We separate two cases according to the invertibility of H_2 .

Case 1.1: H_2 is not pd. Since H_2 is not pd, then by Theorem 2.5, the last column of $\mathcal{H}(\mathcal{G}(t_0, u_0))$ is in the span of the previous ones. But then by rg, the last column of H_2 is in the span of the previous ones. This is the case Theorem 5.1.(2(a)ii).

Case 1.2: H_2 is pd. We separate two cases according to the invertibility of $(\mathcal{H}(A_{\min}))_{\vec{X}(0, k-1)}$.

Case 1.2.1: $\text{rank}(\mathcal{H}(A_{\min}))_{\vec{X}(0, k-1)} = k$. This is the case Theorem 5.1.(2(a)i).

Case 1.2.2: $\text{rank}(\mathcal{H}(A_{\min}))_{\vec{X}(0, k-1)} < k$. We will prove that this case cannot occur. It follows from the assumption in this case that $\text{rank } \mathcal{H}(A_{\min}) = \text{rank } H_2 = k$. Further on, the last column of $\mathcal{H}(A_{\min})$ cannot be in the span of the previous ones (otherwise $\text{rank } \mathcal{H}(A_{\min}) < k$). Hence, by Theorem 2.5, $\mathcal{H}(A_{\min}) = \mathcal{H}(\mathcal{G}(0, 0))$ does not admit a \mathbb{R} -rm. Using this fact and Claim 3, $(0, 0) \in \partial \mathcal{R}_2$. If $t_0 = 0$, then $\mathcal{R}_1 \cap \mathcal{R}_2 = \{(0, 0)\}$, which contradicts to the third condition in (5.11). So $0 < t_0$ must hold. Since $\eta = 0$, Claim 1 implies that $\mathcal{R}_1 = \{(t, 0) : t \geq 0\}$ is a horizontal half-line. By the form of $\partial \mathcal{R}_2$, which is the union of the graphs of two square root functions on the interval $(-\infty, t_0]$, intersecting in the point (t_0, u_0) and such that $(t_0, u_0) \in \partial \mathcal{R}_2$, it follows that $\mathcal{R}_1 \cap \mathcal{R}_2 = \{(0, 0)\}$. Note that by $H_2 \succ 0$, we have $H_2/H_{22} > 0$ and hence $h(t) \not\equiv 0$ (see (5.8)), which implies that the square root functions are indeed not just a horizontal half-line. As above this contradicts to the third condition in (5.11). Hence, Case 1.2.2 cannot occur.

Case 2: $\eta > 0$. By assumptions, $(t_0, u_0) \in \mathcal{R}_1 \cap \mathcal{R}_2$. By Claim 4, $\mathcal{I} \neq \emptyset$ and \mathcal{I} has one or two elements. We separate two cases according to the number of elements in \mathcal{I} .

Case 2.1: \mathcal{I} has two elements. By Claim 4, $H_2/H_{22} > 0$. If H_2 is not pd, then the fact that $\mathcal{H}(\mathcal{G}(t_0, u_0))$ has a \mathbb{R} -rm, implies that $H_2/H_{22} = 0$, which is a contradiction. Indeed, if $H_2/H_{22} > 0$ and H_2 is not pd, then there is a nontrivial column relation among columns X^2, \dots, X^k in H_2 . By Proposition 2.3, the same holds for $\mathcal{H}(\mathcal{G}(t_0, u_0))$. Let $\sum_{i=0}^{k-2} c_i X^{i+2} = \mathbf{0}$ be the nontrivial column relation in $\mathcal{H}(\mathcal{G}(t_0, u_0))$. But then $\mathcal{Z}(x^2 \sum_{i=0}^{k-2} c_i x^i) = \mathcal{Z}(x \sum_{i=0}^{k-2} c_i x^i)$ and it follows by [CF96] that $\sum_{i=0}^{k-2} c_i X^{i+1} = \mathbf{0}$ is also a nontrivial column relation in $\mathcal{H}(\mathcal{G}(t_0, u_0))$. In particular, $H_2/H_{22} = 0$. Hence, H_2 is pd. This is the case Theorem 5.1.(2(b)i).

Case 2.2: \mathcal{I} has one element. Let us denote this element by (\tilde{t}, \tilde{u}) . By Claim 4, $\mathcal{I} = \mathcal{R}_1 \cap \mathcal{R}_2$ or $\partial \mathcal{R}_2 = \mathcal{R}_2 = \{(t, u_0) : t \leq t_0\}$ and $\mathcal{I} \subsetneq \mathcal{R}_1 \cap \mathcal{R}_2 = \{(t, u_0) : \tilde{t} \leq t \leq t_0\}$. We separate two cases according to these two possibilities.

Case 2.2.1: $\mathcal{I} = \mathcal{R}_1 \cap \mathcal{R}_2$. In this case $(t_0, u_0) = (\tilde{t}, \tilde{u})$ and hence $\mathcal{H}(\mathcal{G}(\tilde{t}, \tilde{u}))$ admits a \mathbb{R} -rm. Since $(\tilde{t}, \tilde{u}) \in \partial\mathcal{R}_1$, $\mathcal{H}(\mathcal{G}(\tilde{t}, \tilde{u}))$ is not pd. Hence, by Theorem 2.5, the statement Theorem 5.1.(2(b)ii) holds.

Case 2.2.2: $\partial\mathcal{R}_2 = \mathcal{R}_2 = \{(t, u_0) : t \leq t_0\}$ and $\mathcal{I} \subsetneq \mathcal{R}_1 \cap \mathcal{R}_2 = \{(t, u_0) : \tilde{t} \leq t \leq t_0\}$. By (5.20), it follows that $H_2/H_{22} = 0$ (see the definition (5.8) of $h(\mathbf{t})$). Since H_2 is not pd, Theorem 2.5 used for $\mathcal{H}(\mathcal{G}(t_0, u_0))$, implies that the last column of H_2 is in the span of the others. Hence, the same holds by Proposition 2.3 for $\mathcal{H}(\mathcal{G}(\tilde{t}, \tilde{u}))$ and $\mathcal{H}(\mathcal{G}(\tilde{t}, \tilde{u}))$ admits a \mathbb{R} -rm by Theorem 2.5. Since $\mathcal{H}(\mathcal{G}(\tilde{t}, \tilde{u}))$ is not pd, it in particular satisfies (5.10). Hence, we are in the case Theorem 5.1.(2(b)ii).

This concludes the proof of the implication (5.11) \Rightarrow Theorem 5.1.(2).

Next we prove the implication Theorem 5.1.(2) \Rightarrow (5.11). We separate four cases according to the assumptions in Theorem 5.1.(2).

Case 1: Theorem 5.1.(2(a)i) holds. By Claim 3, $(0, 0) \in \mathcal{R}_1 \cap \mathcal{R}_2$. This and the assumption $\text{rank}(\mathcal{H}(A_{\min}))_{\bar{X}^{(0, k-1)}} = k$, imply by Theorem 2.5, that $\mathcal{H}(\mathcal{G}(0, 0)) = \mathcal{H}(A_{\min})$ admits a \mathbb{R} -rm. This proves (5.11) in case of Theorem 5.1.(2(a)i).

Case 2: Theorem 5.1.(2(a)ii) holds. By Claim 3, $(0, 0) \in \mathcal{R}_1 \cap \mathcal{R}_2$. Since the last column of H_2 is by assumption in the span of the previous ones, the same holds for $\mathcal{H}(\mathcal{G}(0, 0))$ by Proposition 2.3. By Theorem 2.5, $\mathcal{H}(\mathcal{G}(0, 0))$ admits a \mathbb{R} -rm. This proves (5.11) in case of Theorem 5.1.(2(a)ii).

Case 3: Theorem 5.1.(2(b)i) holds. By assumption, $\mathcal{I} = \partial\mathcal{R}_1 \cap \partial\mathcal{R}_2 = \{(t_1, u_1), (t_2, u_2)\}$. Since H_2 is pd, $\partial\mathcal{R}_2$ is not a half-line and hence $u_1 \neq u_2$. By Claim 5, at least one of $\mathcal{H}(\mathcal{G}(t_1, u_1))$ and $\mathcal{H}(\mathcal{G}(t_2, u_2))$ admits a \mathbb{R} -rm. This proves (5.11) in case of Theorem 5.1.(2(b)i).

Case 4: Theorem 5.1.(2(b)ii) holds. The assumptions imply by Theorem 2.5, that $\mathcal{H}(\mathcal{G}(\tilde{t}, \tilde{u}))$ admits a \mathbb{R} -rm. This proves (5.11) in case of Theorem 5.1.(2(b)ii).

This concludes the proof of the implication Theorem 5.1.(2) \Rightarrow (5.11).

Up to now we established the equivalence (1) \Leftrightarrow (2) in Theorem 5.1. It remains to prove the moreover part. We observe again the proof of the implication (2) \Rightarrow (5.11). By Lemma 4.3.(4),

$$(5.26) \quad \text{rank } \widetilde{\mathcal{M}}(k; \beta) = \text{rank } \mathcal{F}(A_{\min}) + \text{rank } \mathcal{H}(A_{\min}).$$

In the proof of the implications Theorem 5.1.(2(a)i) \Rightarrow (5.11) and Theorem 5.1.(2(a)ii) \Rightarrow (5.11) we established that $\mathcal{H}(\mathcal{G}(0, 0))$ has a \mathbb{R} -rm. By Theorem 2.5, there also exists a $(\text{rank } \mathcal{H}(\mathcal{G}(0, 0)))$ -atomic one. By Theorem 2.6, the sequence with the moment matrix $\mathcal{F}(\mathcal{G}(0, 0))$ can be represented by a $(\text{rank } \mathcal{F}(\mathcal{G}(0, 0)))$ -atomic $\mathcal{Z}(ay + x^2 + y^2)$ -rm. By (5.26) and $\mathcal{G}(0, 0) = A_{\min}$ if $\eta = 0$, in these two cases β has a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta))$ -atomic $\mathcal{Z}(p)$ -rm.

In the proof of the implication Theorem 5.1.(2(b)i) \Rightarrow (5.11) we established that $\mathcal{H}(\mathcal{G}(t', u'))$ has a \mathbb{R} -rm for some $(t', u') \in \mathcal{I}$. Analogously as for the point $(0, 0)$ in the previous paragraph, it follows that β has a $(\text{rank } \mathcal{F}(\mathcal{G}(t', u')) + \text{rank } \mathcal{H}(\mathcal{G}(t', u')))$ -atomic $\mathcal{Z}(p)$ -rm. Using (5.13), (5.21) and $\text{rank } H_2 = \text{rank } H_{22} + 1$ (by H_2 being pd), it follows that

$$(5.27) \quad \text{rank } \mathcal{F}(\mathcal{G}(t', u')) + \text{rank } \mathcal{H}(\mathcal{G}(t', u')) = \text{rank } \mathcal{F}(A_{\min}) + \text{rank } H_2 + 1.$$

We separate two cases:

- If $\mathcal{H}(A_{\min})$ is pd, then $\text{rank } \mathcal{H}(A_{\min}) = \text{rank } H_2 + 1$. This, (5.26) and (5.27) imply that β admits a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta))$ -atomic $\mathcal{Z}(p)$ -rm.
- If $\mathcal{H}(A_{\min})$ is not pd, then we must have $\text{rank } \mathcal{H}(A_{\min}) = \text{rank } H_2$. Otherwise we have $(\mathcal{H}(A_{\min}))_{\bar{\mathcal{X}}(1,k)}/H_{22} = 0$ and hence $(\mathcal{H}(A_{\min} - \eta E_{2,2}^{(k+1)}))_{\bar{\mathcal{X}}(1,k)}/H_{22} < 0$, which contradicts to $\mathcal{H}(A_{\min} - \eta E_{2,2}^{(k+1)})$ being psd. Hence, in this case β has a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta) + 1)$ -atomic $\mathcal{Z}(p)$ -rm. Moreover, there cannot exist a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta))$ -atomic $\mathcal{Z}(p)$ -rm. Indeed, since $\eta > 0$, at least $\text{rank } \mathcal{F}(A_{\min}) + 1$ (resp. $\text{rank } H_2$) atoms are needed to represent $\mathcal{F}(\mathcal{G}(t'', u''))$ (resp. $\mathcal{H}(\mathcal{G}(t'', u''))$) for any $(t'', u'') \in \mathcal{R}_1 \cap \mathcal{R}_2$ (see (5.13) and (5.21)). Hence, at least $\text{rank } \mathcal{F}(A_{\min}) + \text{rank } H_2 + 1$ atoms are needed in a $\mathcal{Z}(p)$ -rm for any $(t'', u'') \in \mathcal{R}_1 \cap \mathcal{R}_2$.

In the proof of the implication Theorem 5.1.(2(b)ii) \Rightarrow (5.11) we established that $\mathcal{H}(\mathcal{G}(\tilde{t}, \tilde{u}))$ has a \mathbb{R} -rm. Analogously as for the point $(0, 0)$ in two paragraphs above, it follows that β has a $(\text{rank } \mathcal{F}(\mathcal{G}(\tilde{t}, \tilde{u})) + \text{rank } \mathcal{H}(\mathcal{G}(\tilde{t}, \tilde{u})))$ -atomic $\mathcal{Z}(p)$ -rm. By (5.13) and (5.21), this measure is $(\text{rank } \mathcal{F}(A_{\min}) + \text{rank } H_{22} + 2)$ -atomic.

- If $\mathcal{H}(A_{\min})$ is pd, then $\text{rank } \mathcal{H}(A_{\min}) = \text{rank } H_{22} + 2$. This and (5.26) imply that β admits a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta))$ -atomic $\mathcal{Z}(p)$ -rm.
- If $\mathcal{H}(A_{\min})$ is not pd, then we have $\text{rank } \mathcal{H}(A_{\min}) = \text{rank } H_{22} + 1$, since otherwise the equality $(\mathcal{H}(A_{\min}))_{\bar{\mathcal{X}}(1,k)}/H_{22} = 0$ implies $(\mathcal{H}(A_{\min} - \eta E_{2,2}^{(k+1)}))_{\bar{\mathcal{X}}(1,k)}/H_{22} < 0$, which contradicts to $\mathcal{H}(A_{\min} - \eta E_{2,2}^{(k+1)})$ being psd. Hence, in this case β has a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta) + 1)$ -atomic $\mathcal{Z}(p)$ -rm. Moreover, there cannot exist a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta))$ -atomic $\mathcal{Z}(p)$ -rm in this case. Indeed,

$$(\mathcal{R}_1 \cap \mathcal{R}_2) \setminus \mathcal{I} = (\partial \mathcal{R}_1 \cap \hat{\mathcal{R}}_2) \cup (\hat{\mathcal{R}}_1 \cap \partial \mathcal{R}_2) \cup (\hat{\mathcal{R}}_1 \cap \hat{\mathcal{R}}_2).$$

Using (5.13) and (5.21), in every point from $(\mathcal{R}_1 \cap \mathcal{R}_2) \setminus \mathcal{I}$ at least $\text{rank } \mathcal{F}(A_{\min}) + \text{rank } H_{22} + 2$ atoms are needed in a $\mathcal{Z}(p)$ -rm.

This concludes the proof of the moreover part.

Since for a p -pure sequence with $\widetilde{\mathcal{M}}(k; \beta) \succeq 0$, (5.26) implies that $\mathcal{H}(A_{\min})$ is pd, it follows by the moreover part that the existence of a $\mathcal{Z}(p)$ -rm implies the existence of a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta))$ -atomic $\mathcal{Z}(p)$ -rm. \square

The following example demonstrates the use of Theorem 5.1 to show that there exists a bivariate $y(-2y + x^2 + y^2)$ -pure sequence β of degree 6 with a positive semidefinite $\mathcal{M}(3)$ and without a $\mathcal{Z}(y(-2y + x^2 + y^2))$ -rm.

Example 5.3. Let β be a bivariate degree 6 sequence given by

$$\begin{array}{lll} \beta_{00} = 10, & \beta_{10} = \frac{38}{5}, & \beta_{01} = \frac{39}{5}, \\ \beta_{20} = \frac{602}{25}, & \beta_{11} = \frac{3}{25}, & \beta_{02} = \frac{313}{25}, \\ \beta_{30} = \frac{9152}{125}, & \beta_{21} = \frac{421}{125}, & \beta_{12} = \frac{3}{125}, \\ \beta_{03} = \frac{2709}{125}, & \beta_{40} = \frac{172118}{625}, & \beta_{31} = \frac{27}{625}, \end{array}$$

$$\begin{aligned}
 \beta_{22} &= \frac{2717}{625}, & \beta_{13} &= \frac{3}{625}, & \beta_{04} &= \frac{24373}{625}, \\
 \beta_{50} &= \frac{3303368}{3125}, & \beta_{41} &= \frac{7789}{3125}, & \beta_{32} &= \frac{27}{3125}, \\
 \beta_{23} &= \frac{19381}{3125}, & \beta_{14} &= \frac{3}{3125}, & \beta_{05} &= \frac{224349}{3125}, \\
 \beta_{60} &= 4156, & \beta_{51} &= \frac{243}{15625}, & \beta_{42} &= \frac{44453}{15625}, \\
 \beta_{33} &= \frac{27}{15625}, & \beta_{24} &= \frac{149357}{15625}, & \beta_{15} &= \frac{3}{15625}, \\
 \beta_{06} &= \frac{2094133}{15625}.
 \end{aligned}$$

Assume the notation as in Theorem 5.1. $\widetilde{\mathcal{M}}(3)$ is psd with the eigenvalues ≈ 4445 , ≈ 189.2 , ≈ 16.6 , ≈ 11.9 , ≈ 3.2 , ≈ 1.22 , ≈ 0.57 , ≈ 0.022 , ≈ 0.0030 , 0 and the column relation

$$-2Y^2 + X^2Y + Y^3 = 0.$$

We have that

$$A_{\min} = \begin{pmatrix} \frac{324330}{55873} & \frac{132789}{278915} & \frac{77}{25} & \frac{27}{125} \\ \frac{132789}{278915} & \frac{4180091}{1394575} & \frac{27}{125} & \frac{1493}{625} \\ \frac{77}{25} & \frac{27}{125} & \frac{1493}{625} & \frac{243}{3125} \\ \frac{27}{125} & \frac{1493}{625} & \frac{243}{3125} & \frac{33437}{15625} \end{pmatrix}$$

and so

$$\eta = \frac{77}{25} - \frac{4180091}{1394575} = \frac{4608}{55783}.$$

The matrix H_2 is equal to:

$$H_2 = \begin{pmatrix} 21 & 73 & 273 \\ 73 & 273 & 1057 \\ 273 & 1057 & \frac{64904063}{15625} \end{pmatrix}.$$

The eigenvalues of H_2 are ≈ 4441.1 , ≈ 6.74 , ≈ -0.019 and hence H_2 is not psd. By Theorem 5.1, β does not have a $\mathcal{Z}(y(-2y + x^2 + y^2))$ -rm, since by (2b) of Theorem 5.1, H_2 should be psd.

6. PARABOLIC TYPE RELATION: $p(x, y) = y(x - y^2)$.

In this section we solve the $\mathcal{Z}(p)$ -TMP for the sequence $\beta = \{\beta_i\}_{i,j \in \mathbb{Z}_+, i+j \leq 2k}$ of degree $2k$, $k \geq 3$, where $p(x, y) = y(x - y^2)$. Assume the notation from Section 4. If β admits a $\mathcal{Z}(p)$ -TMP, then $\mathcal{M}(k; \beta)$ must satisfy the relations

$$(6.1) \quad Y^{3+j}X^i = Y^{1+j}X^{i+1} \quad \text{for } i, j \in \mathbb{Z}_+ \text{ such that } i + j \leq k - 3.$$

In the presence of all column relations (6.1), the column space $\mathcal{C}(\mathcal{M}(k; \beta))$ is spanned by the columns in the set

$$(6.2) \quad \mathcal{T} = \vec{X}^{(0,k)} \cup Y\vec{X}^{(0,k-1)} \cup Y^2\vec{X}^{(0,k-2)},$$

where

$$Y^i\vec{X}^{(j,\ell)} := (Y^iX^j, Y^iX^{j+1}, \dots, Y^iX^\ell) \quad \text{with } i, j, \ell \in \mathbb{Z}_+, j \leq \ell, i + \ell \leq k.$$

Let $\widetilde{\mathcal{M}}(k; \beta)$ be as in (4.8). Let

$$(6.3) \quad A_{\min} := A_{12}(A_{22})^\dagger(A_{12})^T.$$

As described in Remark 4.4, A_{\min} might need to be changed to

$$\widehat{A}_{\min} = A_{\min} + \eta(E_{1,k+1}^{(k+1)} + E_{k+1,1}^{(k+1)}),$$

where

$$\eta := (A_{\min})_{2,k} - (A_{\min})_{1,k+1}.$$

Let $\mathcal{F}(\mathbf{A})$ and $\mathcal{H}(\mathbf{A})$ be as in (4.10). Define also the matrix function

$$(6.4) \quad \mathcal{G} : \mathbb{R}^2 \rightarrow S_{k+1}, \quad \mathcal{G}(\mathbf{t}, \mathbf{u}) = \widehat{A}_{\min} + \mathbf{t}E_{1,1}^{(k+1)} + \mathbf{u}E_{k+1,k+1}^{(k+1)}.$$

Write

$$(6.5) \quad \mathcal{H}(\widehat{A}_{\min}) = \begin{matrix} 1 & \vec{X}^{(1,k-1)} & X^k \\ \begin{matrix} 1 \\ \vec{X}^{(1,k-1)T} \end{matrix} & \begin{pmatrix} \beta_{0,0} - (A_{\min})_{1,1} & (h_{12})^T & \beta_{k,0} - (A_{\min})_{2,k} \\ h_{12} & H_{22} & h_{23} \\ \beta_{k,0} - (A_{\min})_{2,k} & (h_{23})^T & \beta_{2k,0} - (A_{\min})_{k+1,k+1} \end{pmatrix} \end{matrix},$$

$$H_1 := (\mathcal{H}(\widehat{A}_{\min}))_{\vec{X}^{(0,k-1)}} = \begin{matrix} 1 & \vec{X}^{(1,k-1)} \\ \begin{matrix} 1 \\ \vec{X}^{(1,k-1)T} \end{matrix} & \begin{pmatrix} \beta_{0,0} - (A_{\min})_{1,1} & (h_{12})^T \\ h_{12} & H_{22} \end{pmatrix} \end{matrix},$$

$$H_2 := (\mathcal{H}(\widehat{A}_{\min}))_{\vec{X}^{(1,k)}} = \begin{matrix} \vec{X}^{(1,k-1)} & X^k \\ \begin{matrix} \vec{X}^{(1,k-1)T} \\ X^k \end{matrix} & \begin{pmatrix} H_{22} & h_{23} \\ (h_{23})^T & \beta_{2k,0} - (A_{\min})_{k+1,k+1} \end{pmatrix} \end{matrix}.$$

Let us define the matrix

$$\begin{aligned} K &:= \mathcal{H}(\widehat{A}_{\min})/H_{22} \\ &= \begin{pmatrix} \beta_{0,0} - (A_{\min})_{1,1} & \beta_{k,0} - (A_{\min})_{2,k} \\ \beta_{k,0} - (A_{\min})_{2,k} & \beta_{2k,0} - (A_{\min})_{k+1,k+1} \end{pmatrix} - \begin{pmatrix} (h_{12})^T \\ (h_{23})^T \end{pmatrix} (H_{22})^\dagger \begin{pmatrix} h_{12} & h_{23} \end{pmatrix} \\ &:= \begin{pmatrix} \beta_{0,0} - (A_{\min})_{1,1} - (h_{12})^T (H_{22})^\dagger h_{12} & \beta_{k,0} - (A_{\min})_{2,k} - (h_{12})^T (H_{22})^\dagger h_{23} \\ \beta_{k,0} - (A_{\min})_{2,k} - (h_{23})^T (H_{22})^\dagger h_{12} & \beta_{2k,0} - (A_{\min})_{k+1,k+1} - (h_{23})^T (H_{22})^\dagger h_{23} \end{pmatrix} \\ &:= \begin{pmatrix} k_{11} & k_{12} \\ k_{12} & k_{22} \end{pmatrix}. \end{aligned}$$

Let

$$\widehat{\mathcal{T}} = \{1, Y, X, XY, X^2, X^2Y, \dots, X^i, X^iY, \dots, X^{k-1}, X^{k-1}Y, X^k\},$$

and

$$(6.6) \quad \widehat{P} \text{ be a permutation matrix such that moment matrix } \widehat{\mathcal{M}}(k; \beta) := \widehat{P}\mathcal{M}(k; \beta)(\widehat{P})^T \text{ has rows and columns indexed in the order } \widehat{\mathcal{T}}, \mathcal{C} \setminus \widehat{\mathcal{T}}.$$

Write

$$(6.7) \quad \widehat{\mathcal{F}}(\mathbf{t}, \mathbf{u}) = \widehat{P}\mathcal{F}(\mathcal{G}(\mathbf{t}, \mathbf{u}))(\widehat{P})^T$$

$$= \begin{pmatrix} 1 & \widehat{\mathcal{T}} \setminus \{1, X^k\} & X^k & \mathcal{C} \setminus \widehat{\mathcal{T}} \\ \widehat{\mathcal{T}} \setminus \{1, X^k\} & (A_{\min})_{1,1} + \mathbf{t} & (f_{12})^T & (A_{\min})_{2,k} & (f_{14})^T \\ X^k & f_{12} & F_{22} & f_{23} & F_{24} \\ \mathcal{C} \setminus \widehat{\mathcal{T}} & (A_{\min})_{2,k} & (f_{23})^T & (A_{\min})_{k+1,k+1} + \mathbf{u} & (f_{34})^T \\ & f_{14} & (F_{24})^T & f_{34} & F_{44} \end{pmatrix}.$$

The solution to the cubic parabolic type relation TMP is the following.

Theorem 6.1. *Let $p(x, y) = y(x - y^2)$ and $\beta := \beta^{(2k)} = (\beta_{i,j})_{i,j \in \mathbb{Z}_+, i+j \leq 2k}$, where $k \geq 3$. Assume also the notation above. Then the following statements are equivalent:*

(1) β has a $\mathcal{Z}(p)$ -representing measure.

(2) $\widetilde{\mathcal{M}}(k; \beta)$ is positive semidefinite, the relations

$$(6.8) \quad \beta_{i,j+3} = \beta_{i+1,j+1} \quad \text{hold for every } i, j \in \mathbb{Z}_+ \text{ with } i + j \leq 2k - 3,$$

$\mathcal{H}(\widehat{A}_{\min})$ is positive semidefinite, defining real numbers

$$(6.9) \quad \begin{aligned} t_1 &= H_1/H_{22} = \beta_{0,0} - (A_{\min})_{1,1} - (h_{12})^T (H_{22})^\dagger h_{12}, \\ u_1 &= H_2/H_{22} = \beta_{2k,0} - (A_{\min})_{k+1,k+1} - (h_{23})^T (H_{22})^\dagger h_{23}, \end{aligned}$$

and the property

$$(6.10) \quad (\mathcal{H}(\widehat{A}_{\min}))_{\overline{\mathcal{X}}(0,k-1)} \succ 0 \quad \text{or} \quad \text{rank}(\mathcal{H}(\widehat{A}_{\min}))_{\overline{\mathcal{X}}(0,k-1)} = \text{rank } \mathcal{H}(\widehat{A}_{\min}),$$

one of the following statements holds:

(a) F_{22} is not positive definite, $\eta = 0$ and (6.10) holds.

(b) F_{22} is positive definite, H_{22} is not positive definite and one of the following holds:

(i) $u_1 = \eta = 0$.

(ii) $u_1 > 0$, $t_1 > 0$, $t_1 u_1 \geq \eta^2$ and $\beta_{k,0} - (A_{\min})_{2,k} = (h_{12})^T (H_{22})^\dagger h_{23}$.

(c) F_{22}, H_{22} are positive definite and one of the following holds:

(i) $\eta = 0$ and (6.10) holds.

(ii) $\eta \neq 0$ and

$$(6.11) \quad (\sqrt{k_{11}k_{22}} - \text{sign}(k_{12})k_{12})^2 \geq \eta^2,$$

where sign is the sign function and $\text{sign}(0) = 0$.

Moreover, if a $\mathcal{Z}(p)$ -representing measure for β exists, then:

- There exists at most $(\text{rank } \widetilde{\mathcal{M}}(k; \beta) + 1)$ -atomic $\mathcal{Z}(p)$ -representing measure.
- There exists a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta))$ -atomic $\mathcal{Z}(p)$ -representing measure if and only if any of the following holds:
 - $\eta = 0$.
 - $\text{rank } \mathcal{H}(A_{\min}) = \text{rank } H_{22} + 2$.
 - $\text{rank } \mathcal{H}(A_{\min}) = \text{rank } H_{22} + 1$ and one of the following holds:
 - * H_{22} is not positive definite and $t_1 u_1 = \eta^2$.
 - * H_{22} is positive definite, $k_{12} = 0$ and $k_{11}k_{22} = \eta^2$.

In particular, a p -pure sequence β with a $\mathcal{Z}(p)$ -representing measure admits a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta))$ -atomic $\mathcal{Z}(p)$ -representing measure.

Remark 6.2. In this remark we explain the idea of the proof of Theorem 6.1 and the meaning of conditions in the statement of the theorem.

By Lemmas 4.1–4.2, the existence of a $\mathcal{Z}(p)$ -rm for β is equivalent to the existence of $t, u \in \mathbb{R}$ such that $\mathcal{F}(\mathcal{G}(t, u))$ admits a $\mathcal{Z}(x - y^2)$ -rm and $\mathcal{H}(\mathcal{G}(t, u))$ admits a \mathbb{R} -rm. Let

$$\mathcal{R}_1 = \{(t, u) \in \mathbb{R}^2 : \mathcal{F}(\mathcal{G}(t, u)) \succeq 0\} \quad \text{and} \quad \mathcal{R}_2 = \{(t, u) \in \mathbb{R}^2 : \mathcal{H}(\mathcal{G}(t, u)) \succeq 0\}.$$


We denote by ∂R_i and $\overset{\circ}{R}_i$ the topological boundary and the interior of the set R_i , respectively. By the necessary conditions for the existence of a $\mathcal{Z}(p)$ -rm [CF04, Fia95, CF96], $\widetilde{\mathcal{M}}(k; \beta)$ must be psd and the relations (6.8) must hold. Then Theorem 6.1.(1) is equivalent to

$$(6.12) \quad \begin{aligned} & \widetilde{\mathcal{M}}(k; \beta) \succeq 0, \text{ the relations (6.8) hold and} \\ & \exists (t_0, u_0) \in \mathcal{R}_1 \cap \mathcal{R}_2 : \mathcal{F}(\mathcal{G}(t_0, u_0)) \text{ and } \mathcal{H}(\mathcal{G}(t_0, u_0)) \text{ admit} \\ & \quad \text{a } \mathcal{Z}(x - y^2)\text{-rm and a } \mathbb{R}\text{-rm, respectively.} \end{aligned}$$

In the proof of Theorem 6.1 we show that (6.12) is equivalent to Theorem 6.1.(2):


(1) First we establish (see Claims 1 and 2 below) that the form of:

- \mathcal{R}_1 is one of the following:



where the left case occurs if $\eta \neq 0$ and the right if $\eta = 0$.

- \mathcal{R}_2 is one of the following:



where the left case occurs if $k_{12} \neq 0$ and the right if $k_{12} = 0$.

(2) If F_{22} is only positive semidefinite but not definite, then we show that (6.12) is equivalent to

$$(6.13) \quad \widetilde{\mathcal{M}}(k; \beta) \succeq 0, \text{ the relations (6.8) hold, } \eta = 0 \text{ and } \mathcal{H}(\mathcal{G}(0, 0)) \text{ admits a } \mathbb{R}\text{-rm.}$$

The latter statement is further equivalent to Theorem 6.1.(2a).

(3) Assume that F_{22} is positive definite and H_{22} is only positive semidefinite but not definite. If:

- $u_1 = 0$, then we show that (6.12) is equivalent to (6.13). The latter statement is further equivalent to Theorem 6.1.(2(b)i).

- $u_1 > 0$, then we show that (6.12) is equivalent to

$$\begin{aligned} & \widetilde{\mathcal{M}}(k; \beta) \succeq 0, \text{ the relations (6.8) hold, } \mathcal{F}(\mathcal{G}(t_1, u_1)) \text{ and} \\ & \mathcal{H}(\mathcal{G}(t_1, u_1)) \text{ admit a } \mathcal{Z}(x - y^2)\text{-rm and a } \mathbb{R}\text{-rm, respectively.} \end{aligned}$$

The latter statement is further equivalent to Theorem 6.1.(2(b)ii).

- $u_1 < 0$, then (6.12) cannot hold.

(4) Assume that F_{22} and H_{22} are positive definite. If:

- $\eta = 0$, then we show that (6.12) is equivalent to (6.13). The latter statement is further equivalent to Theorem 6.1.(2(c)i).

- $\eta \neq 0$, then we show that (6.12) is equivalent to $\mathcal{R}_1 \cap \mathcal{R}_2 \neq \emptyset$. The latter statement is further equivalent to Theorem 6.1.(2(c)ii).

Proof of Theorem 6.1. Let $\mathcal{R}_1, \mathcal{R}_2$ be as in Remark 6.2. As explained in Remark 6.2, Theorem 6.1.(1) is equivalent to (6.12), thus it remains to prove that (6.12) is equivalent to Theorem 6.1.(2).

First we establish a few claims needed in the proof. Claim 1 (resp. 2) describes \mathcal{R}_1 (resp. \mathcal{R}_2) concretely.

Claim 1. Assume that $\widetilde{\mathcal{M}}(k; \beta) \succeq 0$. Then

$$(6.14) \quad \mathcal{R}_1 = \{(t, u) \in \mathbb{R}^2 : t \geq 0, u \geq 0, tu \geq \eta^2\}.$$

If $(t, u) \in \mathcal{R}_1$, we have

$$(6.15) \quad \text{rank } \mathcal{F}(\mathcal{G}(t, u)) = \begin{cases} \text{rank } \mathcal{F}(A_{\min}), & \text{if } \eta = t = u = 0, \\ \text{rank } \mathcal{F}(A_{\min}) + 1, & \text{if } (\eta = t = 0, u > 0) \text{ or } (\eta = u = 0, t > 0) \\ & \text{or } (\eta \neq 0, tu = \eta^2), \\ \text{rank } \mathcal{F}(A_{\min}) + 2, & \text{if } tu > \eta^2. \end{cases}$$

where A_{\min} is as in (6.3).

Proof of Claim 1. Note that

$$(6.16) \quad \begin{aligned} \mathcal{G}(\mathbf{t}, \mathbf{u}) &= A_{\min} + \eta(E_{1,k+1}^{(k+1)} + E_{k+1,1}^{(k+1)}) + \mathbf{t}E_{1,1}^{(k+1)} + \mathbf{u}E_{k+1,k+1}^{(k+1)} \\ &= A_{\min} + \begin{pmatrix} \mathbf{t} & \mathbf{0}_{1,k-1} & \eta \\ \mathbf{0}_{k-1,1} & \mathbf{0}_{k-1} & \mathbf{0}_{k-1,1} \\ \eta & \mathbf{0}_{1,k-1} & \mathbf{u} \end{pmatrix}. \end{aligned}$$

By Lemma 4.3, we have that

$$(6.17) \quad \mathcal{F}(\mathcal{G}(t, u)) \succeq 0 \Leftrightarrow \mathcal{G}(t, u) \succeq A_{\min}$$

Using (6.16), (6.17) and the definition of \mathcal{R}_1 , we have that

$$(6.18) \quad (t, u) \in \mathcal{R}_1 \Leftrightarrow \begin{pmatrix} t & \eta \\ \eta & u \end{pmatrix} \succeq 0 \Leftrightarrow t \geq 0, u \geq 0, tu \geq \eta^2,$$

Define the matrix function

$$(6.23) \quad \begin{aligned} \mathcal{K}(t, \mathbf{u}) &= \mathcal{H}(\mathcal{G}(t, \mathbf{u}))/H_{22} = \mathcal{H}(\widehat{A}_{\min})/H_{22} - \begin{pmatrix} \mathbf{t} & 0 \\ 0 & \mathbf{u} \end{pmatrix} \\ &= K - \begin{pmatrix} \mathbf{t} & 0 \\ 0 & \mathbf{u} \end{pmatrix} = \begin{pmatrix} k_{11} - \mathbf{t} & k_{12} \\ k_{12} & k_{22} - \mathbf{u} \end{pmatrix}. \end{aligned}$$

Claim 2. Assume that $\widetilde{\mathcal{M}}(k; \beta) \succeq 0$. Then

$$(6.24) \quad \begin{aligned} \mathcal{R}_2 &= \{(t, u) \in \mathbb{R}^2 : \mathcal{K}(t, u) \succeq 0\} \\ &= \{(t, u) \in \mathbb{R}^2 : t \leq k_{11}, u \leq k_{22}, (k_{11} - t)(k_{22} - u) \geq k_{12}^2\}. \end{aligned}$$

If $(t, u) \in \mathcal{R}_2$, we have

$$(6.25) \quad \text{rank } \mathcal{H}(\mathcal{G}(t, u)) = \begin{cases} \text{rank } H_{22}, & \text{if } k_{12} = 0, t = k_{11}, u = k_{22}, \\ \text{rank } H_{22} + 1, & \text{if } (k_{11} - t)(k_{22} - u) = k_{12}^2, (t \neq k_{11} \text{ or } u \neq k_{22}), \\ \text{rank } H_{22} + 2, & \text{if } (k_{11} - t)(k_{22} - u) > k_{12}^2. \end{cases}$$

where A_{\min} is as in (6.3).

Proof of Claim 2. Permuting rows and columns of $\mathcal{H}(\mathcal{G}(t, \mathbf{u}))$ we define

$$\widetilde{\mathcal{H}}(\mathcal{G}(t, \mathbf{u})) = \begin{pmatrix} 1 & & X^k & \vec{X}^{(1,k-1)} \\ X^k & \begin{pmatrix} \beta_{0,0} - (A_{\min})_{1,1} - \mathbf{t} & \beta_{k,0} - (A_{\min})_{2,k} \\ \beta_{k,0} - (A_{\min})_{2,k} & \beta_{2k,0} - (A_{\min})_{k+1,k+1} - \mathbf{u} \end{pmatrix} & & \begin{pmatrix} (h_{12})^T \\ (h_{23})^T \end{pmatrix} \\ (\vec{X}^{(1,k-1)})^T & & \begin{pmatrix} h_{12} & h_{23} \end{pmatrix} & H_{22} \end{pmatrix}.$$

Note that

$$\mathcal{H}(\mathcal{G}(t, u)) \succeq 0 \quad \Leftrightarrow \quad \widetilde{\mathcal{H}}(\mathcal{G}(t, u)) \succeq 0$$

and

$$(6.26) \quad \mathcal{H}(A_{\min}) = \begin{pmatrix} 1 & & X^k & \vec{X}^{(1,k-1)} \\ X^k & \begin{pmatrix} \beta_{0,0} - (A_{\min})_{1,1} & \beta_{k,0} - (A_{\min})_{1,k+1} \\ \beta_{k,0} - (A_{\min})_{1,k+1} & \beta_{2k,0} - (A_{\min})_{k+1,k+1} \end{pmatrix} & & \begin{pmatrix} (h_{12})^T \\ (h_{23})^T \end{pmatrix} \\ (\vec{X}^{(1,k-1)})^T & & \begin{pmatrix} h_{12} & h_{23} \end{pmatrix} & H_{22} \end{pmatrix}.$$

By Lemma 4.3.(2), $\mathcal{H}(A_{\min}) \succeq 0$. Permuting rows and columns, this implies that

$$\widetilde{\mathcal{H}}(A_{\min}) = \begin{pmatrix} 1 & & X^k & \vec{X}^{(1,k-1)} \\ X^k & \begin{pmatrix} \beta_{0,0} - (A_{\min})_{1,1} & \beta_{k,0} - (A_{\min})_{1,k+1} \\ \beta_{k,0} - (A_{\min})_{1,k+1} & \beta_{2k,0} - (A_{\min})_{k+1,k+1} \end{pmatrix} & & \begin{pmatrix} (h_{12})^T \\ (h_{23})^T \end{pmatrix} \\ (\vec{X}^{(1,k-1)})^T & & \begin{pmatrix} h_{12} & h_{23} \end{pmatrix} & H_{22} \end{pmatrix} \succeq 0.$$

By Theorem 2.2, used for $(M, C) = (\widetilde{\mathcal{H}}(A_{\min}), H_{22})$, it follows that $H_{22} \succeq 0$ and $h_{12}, h_{23} \in \mathcal{C}(H_{22})$. Let

$$\mathcal{L} : S_2 \rightarrow S_{k+1}, \quad \mathcal{L}(\mathbf{A}) = \begin{pmatrix} \mathbf{A} & \begin{pmatrix} (h_{12})^T \\ (h_{23})^T \end{pmatrix} \\ \begin{pmatrix} h_{12} & h_{23} \end{pmatrix} & H_{22} \end{pmatrix}.$$

be a matrix function. Using Theorem 2.2 again for $(M, C) = (\mathcal{L}(A), H_{22})$, it follows that

$$(6.27) \quad \mathcal{L}(A) \succeq 0 \quad \Leftrightarrow \quad A \succeq \begin{pmatrix} (h_{12})^T \\ (h_{23})^T \end{pmatrix} (H_{22})^\dagger \begin{pmatrix} h_{12} & h_{23} \end{pmatrix}$$

and

$$(6.28) \quad \text{rank } \mathcal{L}(A) = \text{rank } H_{22} + \text{rank} \left(A - \begin{pmatrix} (h_{12})^T \\ (h_{23})^T \end{pmatrix} (H_{22})^\dagger \begin{pmatrix} h_{12} & h_{23} \end{pmatrix} \right)$$

Further, (6.27) implies that

$$\begin{aligned} \tilde{\mathcal{H}}(\mathcal{G}(t, u)) \succeq 0 &\Leftrightarrow \\ \Leftrightarrow \begin{pmatrix} \beta_{0,0} - (A_{\min})_{1,1} - t & \beta_{k,0} - (A_{\min})_{2,k} \\ \beta_{k,0} - (A_{\min})_{2,k} & \beta_{2k,0} - (A_{\min})_{k+1,k+1} - u \end{pmatrix} - \begin{pmatrix} (h_{12})^T \\ (h_{23})^T \end{pmatrix} (H_{22})^\dagger \begin{pmatrix} h_{12} & h_{23} \end{pmatrix} \succeq 0 \\ \Leftrightarrow \mathcal{K}(t, u) \succeq 0, \end{aligned}$$

where we use the definition (6.23) of $\mathcal{K}(t, u)$ in the last equivalence. Moreover, $\text{rank } \tilde{\mathcal{H}}(\mathcal{G}(t, u)) = \text{rank } H_{22} + \text{rank } \mathcal{K}(t, u)$. This proves (6.24) and (6.25). \blacksquare

Claim 3. If $(t, u) \in \mathcal{R}_2 \cap (\mathbb{R}_+)^2$, then

$$tu \leq (\sqrt{k_{11}k_{22}} - \text{sign}(k_{12})k_{12})^2 =: p_{\max}.$$

The equality is achieved if:

- $k_{12} = 0$, in the point $(t, u) = (k_{11}, k_{22})$.
- $k_{12} > 0$, in the point $(t_-, u_-) = (k_{11} - \frac{k_{12}\sqrt{k_{11}}}{\sqrt{k_{22}}}, k_{22} + \frac{k_{12}\sqrt{k_{22}}}{\sqrt{k_{11}}})$.
- $k_{12} < 0$, in the point $(t_+, u_+) = (k_{11} + \frac{k_{12}\sqrt{k_{11}}}{\sqrt{k_{22}}}, k_{22} - \frac{k_{12}\sqrt{k_{22}}}{\sqrt{k_{11}}})$.

Moreover, if $k_{12} \neq 0$, then for every $p \in [0, p_{\max}]$ there exists a point $(\tilde{t}, \tilde{u}) \in \mathcal{R}_2 \cap (\mathbb{R}_+)^2$ such that $\tilde{t}\tilde{u} = p$ and $(k_{11} - \tilde{t})(k_{22} - \tilde{u}) = k_{12}^2$.

Proof of Claim 3. If $k_{12} = 0$, then $(t, u) \in \mathcal{R}_2 \cap (\mathbb{R}_+)^2 = [0, k_{11}] \times [0, k_{22}]$ and Claim 3 is clear.

Assume that $k_{12} \neq 0$. Then clearly tu is maximized in some point satisfying $(k_{11} - t)(k_{22} - u) = k_{12}^2$. Let $f(t) := t(k_{22} - \frac{k_{12}^2}{k_{11} - t})$. We are searching for the maximum of $f(t)$ on the interval $[0, k_{11}]$.

The stationary points of f are $t_\pm = k_{11} \pm \frac{k_{12}\sqrt{k_{11}}}{\sqrt{k_{22}}}$. Then $u_\pm = k_{22} \mp \frac{k_{12}\sqrt{k_{22}}}{\sqrt{k_{11}}}$. If $k_{12} > 0$, then $t_- \in [0, k_{11}]$ (note that $k_{11}k_{22} \geq k_{12}^2$ if $\mathcal{R}_2 \cap (\mathbb{R}_+)^2 \neq \emptyset$). Further on, $t_-u_- = (\sqrt{k_{11}k_{22}} - k_{12})^2$. Similarly, if $k_{12} < 0$, then $t_+ \in [0, k_{11}]$ and $t_+u_+ = (\sqrt{k_{11}k_{22}} + k_{12})^2$. The moreover part follows by noticing that $f(0) = 0$ and hence on the interval $[0, t_\pm]$, f attains all values between 0 and p_{\max} . \blacksquare

In the proof of Theorem 6.1 we will need a few further observations:

- Observe that

$$(6.29) \quad \begin{aligned} (\mathcal{H}(\mathcal{G}(t, u)))_{\vec{X}(0, k-1)} &= H_1 - tE_{1,1}^{(k)}, \\ (\mathcal{H}(\mathcal{G}(t, u)))_{\vec{X}(1, k-1)} &= H_{22}, \\ (\mathcal{H}(\mathcal{G}(t, u)))_{\vec{X}(1, k)} &= H_2 - uE_{k,k}^{(k)}. \end{aligned}$$

- We have

$$(6.30) \quad (\mathcal{H}(\mathcal{G}(t, u)))_{\bar{X}(0, k-1)} / (\mathcal{H}(\mathcal{G}(t, u)))_{\bar{X}(1, k-1)} = H_1/H_{22} - t = t_1 - t,$$

where in the first equality we used (6.29) and in the second the definition of t_1 (see (6.9)).

- We have

$$(6.31) \quad (\mathcal{H}(\mathcal{G}(t, u)))_{\bar{X}(1, k)} / (\mathcal{H}(\mathcal{G}(t, u)))_{\bar{X}(1, k-1)} = H_2/H_{22} - u = u_1 - u,$$

where in the first equality we used (6.29) and in the second the definition of u_1 (see (6.9)).

First we prove the implication (6.12) \Rightarrow Theorem 6.1.(2). By the necessary conditions for the existence of a $\mathcal{Z}(p)$ -rm [CF04, Fia95, CF96], $\widetilde{\mathcal{M}}(k; \beta)$ must be psd and the relations (6.8) must hold. By Lemma 4.3.(2), $\mathcal{F}(A_{\min}) \succeq 0$. Hence,

$$(6.32) \quad \begin{aligned} & \widehat{F} = \widehat{P} \mathcal{F}(A_{\min}) (\widehat{P})^T \\ & = \begin{pmatrix} 1 & & & \\ & (\widehat{\mathcal{T}} \setminus \{1, X^k\})^T & & \\ & & X^k & \\ & & & \mathcal{C} \setminus \widehat{\mathcal{T}} \end{pmatrix} \begin{pmatrix} 1 & \widehat{\mathcal{T}} \setminus \{1, X^k\} & X^k & \mathcal{C} \setminus \widehat{\mathcal{T}} \\ (A_{\min})_{1,1} & (f_{12})^T & (A_{\min})_{1,k+1} & (f_{14})^T \\ f_{12} & F_{22} & f_{23} & F_{24} \\ (A_{\min})_{1,k+1} & (f_{23})^T & (A_{\min})_{k+1,k+1} & (f_{34})^T \\ f_{14} & (F_{24})^T & f_{34} & F_{44} \end{pmatrix} \succeq 0, \end{aligned}$$

where \widehat{P} is as in (6.6). In particular, $F_{22} \succeq 0$. We separate two cases according to the invertibility of F_{22} .

Case 1: F_{22} is not pd. Let $\beta^{(c)}$ be a sequence corresponding to the moment matrix $\mathcal{F}(\mathcal{G}(t_0, u_0))$. Let $\gamma = (\gamma_0, \dots, \gamma_{4k})$ be a sequence defined by $\gamma_i = \beta_{\lfloor \frac{i}{2} \rfloor, i \bmod 2}^{(c)}$. Note that

$$(\widehat{\mathcal{F}}(\mathcal{G}(t_0, u_0)))_{\widehat{\mathcal{T}} \setminus \{1, X^k\}} = (\widehat{F})_{\widehat{\mathcal{T}} \setminus \{1, X^k\}} = F_{22} = A_{\widehat{\gamma}},$$

where $\widehat{\gamma} = (\gamma_2, \dots, \gamma_{4k-2})$. Since F_{22} is not pd, it follows that there is a non-trivial column relation in F_{22} , which is also a column relation in $A_{\widehat{\gamma}}$ by Proposition 2.3. By Theorem 2.7, $\widehat{\gamma}$ has a \mathbb{R} -rm, which implies by Theorem 2.5, that $A_{\widehat{\gamma}}$ is rg. Hence, the last column of $A_{\widehat{\gamma}} = \widehat{\mathcal{F}}(\mathcal{G}(t_0, u_0))$ is in the span of the columns in $\widehat{\mathcal{T}} \setminus \{1, X^k\}$. It follows that

$$(6.33) \quad \begin{pmatrix} (f_{12})^T \\ F_{22} \\ (f_{23})^T \end{pmatrix} (F_{22})^\dagger f_{23} = \begin{pmatrix} (A_{\min})_{2,k} \\ f_{23} \\ (A_{\min})_{k+1,k+1} + u_0 \end{pmatrix}.$$

On the other hand, by construction of \widehat{F} , the column X^k is also in the span of the columns in $\widehat{\mathcal{T}} \setminus \{1, X^k\}$. Hence,

$$(6.34) \quad \begin{pmatrix} (f_{12})^T \\ F_{22} \\ (f_{23})^T \end{pmatrix} (F_{22})^\dagger f_{23} = \begin{pmatrix} (A_{\min})_{1,k+1} \\ f_{23} \\ (A_{\min})_{k+1,k+1} \end{pmatrix}.$$

By (6.33) and (6.34), it follows that $(A_{\min})_{2,k} = (A_{\min})_{1,k+1}$ or equivalently $\eta = 0$, and $u_0 = 0$. Note that

$$(6.35) \quad \begin{aligned} \widehat{\mathcal{F}}(\mathcal{G}(t_0, u_0)) &= \widehat{\mathcal{F}}(\mathcal{G}(t_0, 0)) \succeq \widehat{\mathcal{F}}(\mathcal{G}(0, 0)) = \mathcal{F}(A_{\min}), \\ \mathcal{H}(\mathcal{G}(t_0, u_0)) &= \mathcal{H}(\mathcal{G}(t_0, 0)) \preceq \mathcal{H}(\mathcal{G}(0, 0)) = \mathcal{H}(A_{\min}). \end{aligned}$$

Further on, $\widehat{\mathcal{F}}(A_{\min})$ has a $\mathcal{Z}(x - y^2)$ -rm by Theorem 2.7 and $\mathcal{H}(A_{\min})$ by Theorem 2.5. Indeed, the column X^k of $\widehat{\mathcal{F}}(A_{\min})$ is in the span of the others and since $\mathcal{H}(\mathcal{G}(t_0, 0))$ satisfies the conditions in Theorem 2.5, the same holds for $\mathcal{H}(A_{\min})$. But then the property (6.10) holds (note that $\eta = 0$). This is the case Theorem 6.1.(2a).

Case 2: F_{22} is pd. By Lemma 4.3.(2), $\mathcal{H}(A_{\min}) \succeq 0$ (see (6.26)). In particular, $H_{22} \succeq 0$. We separate two cases according to the invertibility of H_{22} .

Case 2.1: H_{22} is not pd. By (6.31) and Theorem 2.5, it follows that

$$(6.36) \quad u_1 = u_0.$$

By (6.14),

$$(6.37) \quad u_0 \geq 0.$$

We separate two cases according to the value of u_1 .

Case 2.1.1: $u_1 = 0$. By (6.36), it follows that $u_0 = 0$. Note that

$$(6.38) \quad (\widehat{\mathcal{F}}(\mathcal{G}(t_0, u_0)))_{\widehat{\mathcal{T}} \setminus \{1\}} = (\widehat{\mathcal{F}}(\mathcal{G}(t_0, 0)))_{\widehat{\mathcal{T}} \setminus \{1\}} = (\widehat{F})_{\widehat{\mathcal{T}} \setminus \{1\}}.$$

Since in \widehat{F} we have the column relation (6.34) by construction, (6.38) and Proposition 2.3 imply that

$$(\widehat{\mathcal{F}}(\mathcal{G}(t_0, 0)))_{\widehat{\mathcal{T}}, \widehat{\mathcal{T}} \setminus \{1, X^k\}} (F_{22})^\dagger f_{23} = (\widehat{\mathcal{F}}(\mathcal{G}(t_0, 0)))_{\widehat{\mathcal{T}}, \{X^k\}},$$

or equivalently (6.33) with $u_0 = 0$. By (6.33) and (6.34), it follows that $(A_{\min})_{2,k} = (A_{\min})_{1,k+1}$ or equivalently $\eta = 0$. This is the case Theorem 6.1.(2(b)i).

Case 2.1.2: $u_1 > 0$. Since the column X^k of $\mathcal{H}(\mathcal{G}(t_0, u_1))$ is in the span of the columns in $\vec{X}^{(1,k-1)}$, it first follows by observing the first row of $\mathcal{H}(\mathcal{G}(t_0, u_1))$ that

$$(6.39) \quad \beta_{k,0} - (A_{\min})_{2,k} = (h_{12})^T (H_{22})^\dagger h_{23}.$$

Further on,

$$(6.40) \quad \mathcal{H}(\mathcal{G}(t, u_1)) / (\mathcal{H}(\mathcal{G}(t, u_1)))_{\vec{X}^{(1,k)}} = (\mathcal{H}(\mathcal{G}(t, u_1)))_{\vec{X}^{(0,k-1)}} / (\mathcal{H}(\mathcal{G}(t, u_1)))_{\vec{X}^{(1,k-1)}} = t_1 - t,$$

where we used (6.30) in the second equality. By (6.40) and Theorem 2.2 used for $(M, C) = (\mathcal{H}(\mathcal{G}(t, u_1)), (\mathcal{H}(\mathcal{G}(t, u_1)))_{\vec{X}^{(1,k)}})$, it follows that $\mathcal{H}(\mathcal{G}(t_1, u_1)) \succeq 0$. By Theorem 2.5, $\mathcal{H}(\mathcal{G}(t_1, u_1))$ admits a \mathbb{R} -rm. Note that

$$(6.41) \quad \widehat{\mathcal{F}}(\mathcal{G}(t_0, u_0)) = \widehat{\mathcal{F}}(\mathcal{G}(t_0, u_1)) \preceq \widehat{\mathcal{F}}(\mathcal{G}(t_1, u_1)),$$

where we used that $t_0 \leq t_1$ by (6.40). By Theorem 2.7, $(\widehat{\mathcal{F}}(\mathcal{G}(t_1, u_1)))_{\widehat{\mathcal{T}} \setminus \{X^k\}}$ must be pd. (Here we used that since $u_1 > 0$ and $F_{22} \succ 0$, it follows that $(\widehat{\mathcal{F}}(\mathcal{G}(t_1, u_1)))_{\widehat{\mathcal{T}} \setminus \{1\}} \succ 0$.) Therefore Claim 1 implies that $t_1 > 0$ and $t_1 u_1 \geq \eta^2$. Together with (6.39), this is the case Theorem 6.1.(2(b)ii).

Case 2.2: H_{22} is pd. We separate two cases according to the value of η .

Case 2.2.1: $\eta = 0$. By Lemma 4.3.(2), $\mathcal{H}(A_{\min}) \succeq 0$ (see (6.26)).

If $\mathcal{H}(A_{\min})$ does not admit a \mathbb{R} -rm, it follows by Theorem 2.5, that $(\mathcal{H}(A_{\min}))_{\vec{X}^{(0,k-1)}}$ is not pd and $u_1 > 0$. Equivalently,

$$t_1 = (\mathcal{H}(A_{\min}))_{\vec{X}^{(0,k-1)}} / H_{22} = 0,$$

which by (6.30) implies that $t_0 = 0$. By Theorem 2.7, since $\widehat{\mathcal{F}}(\mathcal{G}(t_0, u_0)) = \widehat{\mathcal{F}}(\mathcal{G}(0, u_0))$ admits a $\mathcal{Z}(x - y^2)$ -rm, $F_{22} \succ 0$ and $(\widehat{\mathcal{F}}(\mathcal{G}(0, u_0)))_{\widehat{\mathcal{T}} \setminus \{X^k\}}$ is not pd, it follows that $u_0 = 0$. But then $\mathcal{H}(\mathcal{G}(t_0, u_0)) = \mathcal{H}(\mathcal{G}(0, 0)) = \mathcal{H}(A_{\min})$ does not admit a \mathbb{R} -rm, which is a contradiction.

Hence, $\mathcal{H}(A_{\min})$ admits a \mathbb{R} -rm, which is equivalent to (6.10) (using $\eta = 0$). This is the case Theorem 6.1.(2(c)i).

Case 2.2.2: $\eta \neq 0$. By (6.15) it follows that $t_0 u_0 \geq \eta^2$. This fact and Claim 3 imply the second condition in the case Theorem 6.1.(2(c)ii).

This concludes the proof of the implication (6.12) \Rightarrow Theorem 6.1.(2).

Next we prove the implication Theorem 6.1.(2) \Rightarrow (6.12). We separate five cases according to the assumptions in Theorem 6.1.(2).

Case 1: Theorem 6.1.(2a) holds. By Lemma 4.3.(2), $\mathcal{F}(A_{\min}) \succeq 0$ and $\mathcal{H}(A_{\min}) \succeq 0$. Since $\eta = 0$, both matrices have a moment structure. Since by construction, the column X^k of $\widehat{\mathcal{F}}(A_{\min})$ is in the span of the others, it has a $\mathcal{Z}(x - y^2)$ -rm by Theorem 2.7. Since $\mathcal{H}(A_{\min})$ satisfies (6.10) (using $\eta = 0$), it admits a \mathbb{R} -rm by Theorem 2.5. This proves (6.12) in this case.

Case 2: Theorem 6.1.(2(b)i) holds. By the same reasoning as in the Case 1 above, $\widehat{\mathcal{F}}(A_{\min})$ has a $\mathcal{Z}(x - y^2)$ -rm. Since $u_1 = 0$, the column X^k of $\mathcal{H}(A_{\min})$ is in the span of the other columns. By Theorem 2.5, $\mathcal{H}(A_{\min})$ admits a \mathbb{R} -rm. This proves (6.12) in this case.

Case 3: Theorem 6.1.(2(b)ii) holds. By (6.30), (6.31) and the fourth assumption of (2(b)ii), it follows that $\mathcal{H}(\mathcal{G}(t_1, u_1))$ is psd and the columns $1, X^k$ are in the span of the columns in $\vec{X}^{(1,k-1)}$. By Theorem 2.5, $\mathcal{H}(\mathcal{G}(t_1, u_1))$ admits a \mathbb{R} -rm. Since $(t_1, u_1) \in \mathcal{R}_1$ by (6.14) and the assumptions in (2(b)ii), it follows that $\widehat{\mathcal{F}}(\mathcal{G}(t_1, u_1))$ is psd and by construction, $(\widehat{\mathcal{F}}(\mathcal{G}(t_1, u_1)))_{\widehat{\mathcal{T}} \setminus \{X^k\}}$ is pd. By Theorem 2.7, it has a $\mathcal{Z}(x - y^2)$ -rm. This proves (6.12) in this case.

Case 4: Theorem 6.1.(2(c)i) holds. $\widehat{\mathcal{F}}(A_{\min})$ has a $\mathcal{Z}(x - y^2)$ -rm and $\mathcal{H}(A_{\min})$ has a \mathbb{R} -rm by the same reasoning as in the Case 1 above. This proves (6.12) in this case.

Case 5: Theorem 6.1.(2(c)ii) holds. We separate three cases according to the sign of k_{12} .

- If $k_{12} = 0$, then by Claim 2, $\mathcal{H}(\mathcal{G}(k_{11}, k_{22}))$ is psd and the column X^k is in the span of the previous ones. Since $\mathcal{H}(\mathcal{G}(0, 0)) = \mathcal{H}(\widehat{A}_{\min})$ is psd by assumption, it follows that $k_{11} \geq 0$ and $k_{22} \geq 0$. Since $\eta \neq 0$ and $k_{11} k_{22} \geq \eta^2$ by (6.11), it follows that $k_{11} > 0$ and $k_{22} > 0$. By Claim 1, $\widehat{\mathcal{F}}(\mathcal{G}(k_{11}, k_{22})) \succ 0$. By Theorem 2.7, it has a $\mathcal{Z}(x - y^2)$ -rm. This proves (6.12) in this case.
- If $k_{12} > 0$, then by Claim 3, $\mathcal{H}(\mathcal{G}(t_-, u_-))$ is psd and $t_- u_- \geq \eta^2$. By construction, $\text{rank } \mathcal{H}(\mathcal{G}(t_-, u_-)) = k$ and since $t_- < k_{11}$, it follows that $(\mathcal{H}(\mathcal{G}(t_-, u_-)))_{\vec{X}^{(0,k-1)}}$ is pd. Hence, the column X^k of $\mathcal{H}(\mathcal{G}(t_-, u_-))$ is in the span of the others. By Theorem 2.5, $\mathcal{H}(\mathcal{G}(t_-, u_-))$ admits a \mathbb{R} -rm. By Claim 1 and $t_- u_- \geq \eta^2$, it follows that $\widehat{\mathcal{F}}(\mathcal{G}(t_-, u_-)) \succeq$

0. Since $t_- > 0$, it follows that $(\widehat{F}(\mathcal{G}(t_-, u_-)))_{\widehat{\mathcal{T}} \setminus \{X^k\}}$ is pd. By Theorem 2.7, it has a $\mathcal{Z}(x - y^2)$ -rm. This proves (6.12) in this case.
- If $k_{12} < 0$, then the proof of (6.12) is analogous to the case $k_{12} > 0$ by replacing (t_-, u_-) with (t_+, u_+) .

This concludes the proof of the implication Theorem 6.1.(2) \Rightarrow (6.12).

By now we established the equivalence (1) \Leftrightarrow (2) in Theorem 6.1. It remains to prove the moreover part. We observe again the proof of the implication (2) \Rightarrow (6.12). By Lemma 4.3.(4),

$$(6.42) \quad \text{rank } \widetilde{\mathcal{M}}(k; \beta) = \text{rank } \widehat{\mathcal{F}}(A_{\min}) + \text{rank } \mathcal{H}(A_{\min}).$$

In the proofs of the implications Theorem 6.1.(2a) \Rightarrow (6.12), Theorem 6.1.(2(b)i) \Rightarrow (6.12) and Theorem 6.1.(2(c)i) \Rightarrow (6.12), we established that $\widehat{\mathcal{F}}(A_{\min})$ and $\mathcal{H}(A_{\min})$ admit a $\mathcal{Z}(x - y^2)$ -rm and a \mathbb{R} -rm, respectively. By Theorems 2.5 and 2.7, there also exist a $(\text{rank } \widehat{\mathcal{F}}(A_{\min}))$ -atomic and a $(\text{rank } \mathcal{H}(A_{\min}))$ -atomic rms. By (6.42), β has a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta))$ -atomic $\mathcal{Z}(p)$ -rm.

Assume that Theorem 6.1.(2(b)ii) holds. We separate two cases according to the value of η :

- $\eta = 0$. We separate two cases according to the existence of a \mathbb{R} -rm of $\mathcal{H}(A_{\min})$:
 - *The last column of $\mathcal{H}(A_{\min})$ is in the span of the previous ones.* Then as in the previous paragraph, $\widehat{\mathcal{F}}(A_{\min})$ and $\mathcal{H}(A_{\min})$ admit a $(\text{rank } \widehat{\mathcal{F}}(A_{\min}))$ -atomic $\mathcal{Z}(x - y^2)$ -rm and a $(\text{rank } \mathcal{H}(A_{\min}))$ -atomic \mathbb{R} -rm, respectively. Hence, β has a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta))$ -atomic $\mathcal{Z}(p)$ -rm.
 - *The last column of $\mathcal{H}(A_{\min})$ is not in the span of the previous ones.* Since also $t_1 > 0$, it follows that $\text{rank } \mathcal{H}(A_{\min}) = \text{rank } H_{22} + 2$. But then $\text{rank } \mathcal{H}(\mathcal{G}(t_1, u_1)) = \text{rank } H_{22}$ and $\text{rank } \widehat{\mathcal{F}}(\mathcal{G}(t_1, u_1)) = \text{rank } \widehat{\mathcal{F}}(A_{\min}) + 2$ (see (6.15)). This implies that $\widetilde{\mathcal{M}}(\beta; k)$ admits a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta))$ -atomic $\mathcal{Z}(p)$ -rm.
- $\eta \neq 0$. We separate two cases according to $\text{rank } \mathcal{H}(A_{\min})$, which can be either $\text{rank } H_{22} + 2$ or $\text{rank } H_{22} + 1$ (since $t_1 > 0$).
 - $\text{rank } \mathcal{H}(A_{\min}) = \text{rank } H_{22} + 2$. Then as in the second Case of the case $\eta = 0$ above, in the point (t_1, u_1) there is a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta))$ -atomic $\mathcal{Z}(p)$ -rm for β . (Note that $t_1 u_1$ is automatically strictly larger than η^2 , otherwise the measure was $(\text{rank } \widetilde{\mathcal{M}}(k; \beta) - 1)$ -atomic, which is not possible.)
 - $\text{rank } \mathcal{H}(A_{\min}) = \text{rank } H_{22} + 1$. In this case we have

$$\begin{aligned} \text{rank } \mathcal{H}(\mathcal{G}(t_1, u_1)) + \text{rank } \widehat{\mathcal{F}}(\mathcal{G}(t_1, u_1)) &= \text{rank } H_{22} + \text{rank } \widehat{\mathcal{F}}(\mathcal{G}(t_1, u_1)) \\ &= \begin{cases} \text{rank } H_{22} + \text{rank } \widehat{\mathcal{F}}(A_{\min}) + 1, & \text{if } t_1 u_1 = \eta^2, \\ \text{rank } H_{22} + \text{rank } \widehat{\mathcal{F}}(A_{\min}) + 2, & \text{if } t_1 u_1 > \eta^2, \end{cases} \\ &= \begin{cases} \text{rank } \widetilde{\mathcal{M}}(k; \beta), & \text{if } t_1 u_1 = \eta^2, \\ \text{rank } \widetilde{\mathcal{M}}(k; \beta) + 1, & \text{if } t_1 u_1 > \eta^2, \end{cases} \end{aligned}$$

where we used (6.15) in the second and (6.42) in the third equality. Hence, β has a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta))$ -atomic rm if $t_1 u_1 = \eta^2$ and $(\text{rank } \widetilde{\mathcal{M}}(k; \beta) + 1)$ -atomic rm if $t_1 u_1 > \eta^2$. It remains to show that in the case $t_1 u_1 > \eta^2$, there does not exist a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta))$ -atomic rm. Since H_{22} is not pd and $u_1 > 0$, if $\mathcal{H}(\mathcal{G}(t', u'))$ has a

\mathbb{R} -rm, then $u' = u_1$. Since $\eta \neq 0$, then $\widehat{\mathcal{F}}(\mathcal{G}(t', u_1))$ with a $\mathcal{Z}(x - y^2)$ -rm is at least $(\text{rank } \widehat{\mathcal{F}}(A_{\min}) + 1)$ -atomic (see (6.15)). If $t' \neq t_1$, then $\text{rank } \mathcal{H}(\mathcal{G}(t', u_1)) = \text{rank } H_{22} + 1$. Hence,

$$\text{rank } \mathcal{H}(\mathcal{G}(t', u_1)) + \text{rank } \widehat{\mathcal{F}}(\mathcal{G}(t', u_1)) \geq (\text{rank } H_{22} + 1) + (\text{rank } \widehat{\mathcal{F}}(A_{\min}) + 1) = \text{rank } \widetilde{\mathcal{M}}(k; \beta) + 1,$$

where we used (6.42) in the last equality.

- Assume that Theorem 6.1.(2(c)iii) holds. We separate two cases according to the value of k_{12} .

- $k_{12} = 0$. We separate two cases according to $\text{rank } \mathcal{H}(A_{\min})$, i.e., $\text{rank } \mathcal{H}(A_{\min}) \in \{k, k + 1\}$. Note that $\text{rank } \mathcal{H}(A_{\min})$ cannot be $k - 1$, since $\eta \neq 0$ and $k_{12} = 0$ imply that $(\mathcal{H}(A_{\min})/H_{22})_{12} \neq 0$.

- * $\text{rank } \mathcal{H}(A_{\min}) = k + 1$. Then as in the second case of the case $\eta = 0$ of Theorem 6.1.(2(b)ii) above, in the point (t_1, u_1) there is a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta))$ -atomic $\mathcal{Z}(p)$ -rm for β . (Note that $t_1 u_1$ is automatically strictly larger than η^2 , otherwise the measure was $(\text{rank } \widetilde{\mathcal{M}}(k; \beta) - 1)$ -atomic, which is not possible.)

- * $\text{rank } \mathcal{H}(A_{\min}) = k$. In this case we have

$$\begin{aligned} \text{rank } \mathcal{H}(\mathcal{G}(k_{11}, k_{22})) + \text{rank } \mathcal{F}(\mathcal{G}(k_{11}, k_{22})) &= \begin{cases} \text{rank } H_{22} + \text{rank } \mathcal{F}(A_{\min}) + 1, & \text{if } k_{11}k_{22} = \eta^2, \\ \text{rank } H_{22} + \text{rank } \mathcal{F}(A_{\min}) + 2, & \text{if } k_{11}k_{22} > \eta^2, \end{cases} \\ &= \begin{cases} \text{rank } \widetilde{\mathcal{M}}(k; \beta), & \text{if } k_{11}k_{22} = \eta^2, \\ \text{rank } \widetilde{\mathcal{M}}(k; \beta) + 1, & \text{if } k_{11}k_{22} > \eta^2, \end{cases} \end{aligned}$$

where we used (6.15) in the first and (6.42) in the second equality. Hence, β has a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta))$ -atomic rm if $k_{11}k_{22} = \eta^2$ and $(\text{rank } \widetilde{\mathcal{M}}(k; \beta) + 1)$ -atomic rm if $k_{11}k_{22} > \eta^2$. It remains to show that in the case $k_{11}k_{22} > \eta^2$, there does not exist a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta))$ -atomic rm. Since $\eta \neq 0$, if $\mathcal{F}(\mathcal{G}(t', u'))$ is psd, it follows that $t'u' \geq \eta^2$ by (6.14). But then if $\widehat{\mathcal{F}}(\mathcal{G}(t', u'))$ also admits a $\mathcal{Z}(x - y^2)$ -rm, this rm is at least $(\text{rank } \widehat{\mathcal{F}}(A_{\min}) + 1)$ -atomic (see (6.15)). If $t' < k_{11}$ or $u' < k_{22}$, then $\text{rank } \mathcal{H}(\mathcal{G}(t', u')) \geq \text{rank } H_{22} + 1$. Hence,

$$\text{rank } \mathcal{H}(\mathcal{G}(t', u')) + \text{rank } \widehat{\mathcal{F}}(\mathcal{G}(t', u')) \geq (\text{rank } H_{22} + 1) + (\text{rank } \widehat{\mathcal{F}}(A_{\min}) + 1) = \text{rank } \widetilde{\mathcal{M}}(k; \beta) + 1,$$

where we used (6.42) in the last equality.

- $k_{12} \neq 0$. We separate two cases according to $\text{rank } \mathcal{H}(A_{\min})$, i.e., $\text{rank } \mathcal{H}(A_{\min}) \in \{k, k + 1\}$. Note that $\text{rank } \mathcal{H}(A_{\min})$ cannot be $k - 1$, since otherwise $\mathcal{H}(\widehat{A}_{\min})/H_{22} = \begin{pmatrix} 0 & \eta \\ \eta & 0 \end{pmatrix}$, which cannot be psd by $\eta \neq 0$. By Claim 3, there is a point $(\tilde{t}, \tilde{u}) \in \mathcal{R}_2 \cap (\mathbb{R}_+)^2$, such that $\tilde{t}\tilde{u} = \eta^2$ and $(k_{11} - \tilde{t})(k_{22} - \tilde{u}) = k_{12}^2$. By (6.15) and (6.25) we have

$$\begin{aligned} \text{rank } \mathcal{H}(\mathcal{G}(\tilde{t}, \tilde{u})) + \text{rank } \widehat{\mathcal{F}}(\mathcal{G}(\tilde{t}, \tilde{u})) &= (\text{rank } H_{22} + 1) + (\text{rank } \widehat{\mathcal{F}}(A_{\min}) + 1) \\ &= \begin{cases} \text{rank } \widetilde{\mathcal{M}}(k; \beta), & \text{if } \text{rank } \mathcal{H}(A_{\min}) = k + 1, \\ \text{rank } \widetilde{\mathcal{M}}(k; \beta) + 1, & \text{if } \text{rank } \mathcal{H}(A_{\min}) = k, \end{cases} \end{aligned}$$

where we used (6.42) in the second equality. It remains to show that in the case $\text{rank } \mathcal{H}(A_{\min}) = k$, there does not exist a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta))$ -atomic rm. Since $\eta \neq 0$,

if $\widehat{\mathcal{F}}(\mathcal{G}(t', u'))$ is psd, it follows that $t'u' \geq \eta^2$ by (6.14). But then if $\widehat{\mathcal{F}}(\mathcal{G}(t', u'))$ also admits a $\mathcal{Z}(x - y^2)$ -rm, this rm is at least $(\text{rank } \widehat{\mathcal{F}}(A_{\min}) + 1)$ -atomic (see (6.15)). Since $k_{12} \neq 0$, $\text{rank } \mathcal{H}(\mathcal{G}(t', u')) \geq \text{rank } H_{22} + 1$ by (6.25). Hence,

$$\text{rank } \mathcal{H}(\mathcal{G}(t', u')) + \text{rank } \widehat{\mathcal{F}}(\mathcal{G}(t', u')) \geq (\text{rank } H_{22} + 1) + (\text{rank } \widehat{\mathcal{F}}(A_{\min}) + 1) = \text{rank } \widetilde{\mathcal{M}}(k; \beta) + 1,$$

where we used (6.42) in the last equality.

This concludes the proof of the moreover part.

Since for a p -pure sequence with $\widetilde{\mathcal{M}}(k; \beta) \succeq 0$, (6.42) implies that $\mathcal{H}(A_{\min})$ is pd, it follows by the moreover part that the existence of a $\mathcal{Z}(p)$ -rm implies the existence of a $(\text{rank } \widetilde{\mathcal{M}}(k; \beta))$ -atomic $\mathcal{Z}(p)$ -rm. \square

The following example demonstrates the use of Theorem 6.1 to show that there exists a bivariate $y(x - y^2)$ -pure sequence β of degree 6 with a positive semidefinite $\mathcal{M}(3)$ and without a $\mathcal{Z}(y(x - y^2))$ -rm.

Example 6.3. Let β be a bivariate degree 6 sequence given by

$$\begin{aligned} \beta_{00} &= \frac{1228153}{1372615}, & \beta_{10} &= \frac{97}{10}, & \beta_{01} &= \frac{21}{10}, \\ \beta_{20} &= \frac{2289}{10}, & \beta_{11} &= \frac{441}{10}, & \beta_{02} &= \frac{91}{10}, \\ \beta_{30} &= \frac{67207}{10}, & \beta_{21} &= \frac{12201}{10}, & \beta_{12} &= \frac{455}{2}, \\ \beta_{03} &= \frac{441}{10}, & \beta_{40} &= \frac{2142693}{10}, & \beta_{31} &= \frac{376761}{10}, \\ \beta_{22} &= \frac{67171}{10}, & \beta_{13} &= \frac{12201}{10}, & \beta_{04} &= \frac{455}{2}, \\ \beta_{50} &= \frac{71340727}{10}, & \beta_{41} &= \frac{12313161}{10}, & \beta_{32} &= \frac{428519}{2}, \\ \beta_{23} &= \frac{376761}{10}, & \beta_{14} &= \frac{67171}{10}, & \beta_{05} &= \frac{12201}{10}, \\ \beta_{60} &= \frac{2438236509}{10}, & \beta_{51} &= \frac{415998681}{10}, & \beta_{42} &= \frac{71340451}{10}, \\ \beta_{33} &= \frac{12313161}{10}, & \beta_{24} &= \frac{428519}{2}, & \beta_{15} &= \frac{376761}{10}, \\ \beta_{06} &= \frac{67171}{10}. \end{aligned}$$

Assume the notation as in Theorem 6.1. $\widetilde{\mathcal{M}}(3)$ is psd with the eigenvalues $\approx 2.51 \cdot 10^8$, ≈ 47179 , ≈ 112.1 , ≈ 7.4 , ≈ 1.11 , ≈ 0.1 , ≈ 0.03 , ≈ 0.0005 , $\approx 4.9 \cdot 10^{-6}$, 0, and the column relation $Y^3 = YX$. We have that

$$A_{\min} = \begin{pmatrix} \frac{5537}{9230} & \frac{91}{10} & \frac{455}{2} & \frac{61999553}{9230} \\ \frac{91}{10} & \frac{455}{2} & \frac{67171}{10} & \frac{428519}{2} \\ \frac{455}{2} & \frac{67171}{10} & \frac{428519}{2} & \frac{71340451}{10} \\ \frac{61999553}{9230} & \frac{428519}{2} & \frac{71340451}{10} & \frac{450098209309}{1846} \end{pmatrix}$$

and so

$$\eta = \frac{67171}{10} - \frac{61999553}{9230} = -\frac{72}{923}.$$

The matrices F_{22} and H_{22} are equal to:

$$F_{22} = \begin{pmatrix} \frac{91}{10} & \frac{441}{10} & \frac{455}{2} & \frac{12201}{10} & \frac{67171}{10} \\ \frac{441}{10} & \frac{455}{2} & \frac{12201}{10} & \frac{67171}{10} & \frac{376761}{10} \\ \frac{455}{2} & \frac{12201}{10} & \frac{67171}{10} & \frac{376761}{10} & \frac{428519}{2} \\ \frac{12201}{10} & \frac{67171}{10} & \frac{376761}{10} & \frac{428519}{2} & \frac{12313161}{10} \\ \frac{67171}{10} & \frac{376761}{10} & \frac{428519}{2} & \frac{12313161}{10} & \frac{71340451}{10} \end{pmatrix}, \quad H_{22} = \begin{pmatrix} \frac{7}{5} & \frac{18}{5} \\ \frac{18}{5} & \frac{49}{5} \end{pmatrix}.$$

They are both pd with the eigenvalues $\approx 7.3 \cdot 10^6$, ≈ 1987.6 , ≈ 5.6 , ≈ 0.099 , ≈ 0.0013 and ≈ 11.1 , ≈ 0.068 , respectively. The matrix K is equal to

$$K = \begin{pmatrix} k_{11} & k_{12} \\ k_{12} & k_{22} \end{pmatrix} = \begin{pmatrix} \frac{6050329}{48143098510} & \frac{3}{95} \\ \frac{3}{95} & \frac{4941414}{87685} \end{pmatrix}$$

and thus

$$(6.43) \quad (\sqrt{k_{11}k_{12}} - k_{12})^2 - \eta^2 = -0.0033 < 0.$$

By Theorem 6.1, β does not have a $\mathcal{Z}(y(x - y^2))$ -rm, since by (2(c)ii) of Theorem 6.1, (6.43) should be positive.

REFERENCES

- [Alb69] A. Albert, *Conditions for positive and nonnegative definiteness in terms of pseudoinverses*, SIAM J. Appl. Math. **17** (1969), 434–440.
- [Akh65] N.I. Akhiezer. *The classical moment problem and some related questions in analysis*, New York: Hafner Publishing Co., 1965.
- [AK62] N.I. Akhiezer, M. Krein. *Some questions in the theory of moments*, Transl. Math. Monographs 2. Providence: American Math. Soc., 1962.
- [BZ21] A. Bhardwaj, A. Zalar. *The tracial moment problem on quadratic varieties*, J. Math. Anal. Appl. **498** (2021). Available from: <https://doi.org/10.1016/j.jmaa.2021.124936>.
- [BW11] M. Bakonyi, H.J. Woerdeman, *Matrix Completions, Moments, and Sums of Hermitian Squares*, Princeton University Press, Princeton, 2011.
- [Ble15] G. Blekherman. *Positive Gorenstein ideals*, Proc. Amer. Math. Soc. **143** (2015) 69–86. Available from: <https://doi.org/10.1090/S0002-9939-2014-12253-2>.
- [BF20] G. Blekherman, L. Fialkow. *The core variety and representing measures in the truncated moment problem*, Journal of Operator Theory **84** (2020) 185–209.
- [CGIK+] R. Curto, M. Ghasemi, M. Infusino, S. Kuhlmann. *The truncated moment problems for unital commutative \mathbb{R} -algebras*, Journal of Operator Theory, to appear. Available from: <https://arxiv.org/pdf/2009.05115.pdf>.
- [CHM74] D. Carlson, E. Haynsworth, T. Markham, *A generalization of the Schur complement by means of the Moore-Penrose inverse*, SIAM J. Appl. Math. **26**(1) (1974) 169–175.
- [CH69] D. Crabtree, E. Haynsworth, *An identity for the Schur complement of a matrix*. Proc. Am. Math. Soc. **22** (1969) 364–366.
- [CF91] R. Curto, L. Fialkow, *Recursiveness, positivity, and truncated moment problems*, Houston J. Math. **17** (1991) 603–635.
- [CF96] R. Curto, L. Fialkow, *Solution of the truncated complex moment problem for flat data*, Mem. Amer. Math. Soc. **119** (1996).
- [CF02] R. Curto, L. Fialkow, *Solution of the singular quartic moment problem*, J. Operator Theory **48** (2002) 315–354.

- [CF04] R. Curto, L. Fialkow, *Solution of the truncated parabolic moment problem*, Integral Equations Operator Theory **50** (2004), 169–196.
- [CF05] R. Curto, L. Fialkow, *Solution of the truncated hyperbolic moment problem*, Integral Equations Operator Theory **52** (2005) 181–218.
- [CF05b] R. Curto, L. Fialkow, *Truncated K -moment problems in several variables*, J. Operator Theory **54** (2005) 189–226.
- [CF08] R. Curto, L. Fialkow, *An analogue of the Riesz-Haviland theorem for the truncated moment problem*, J. Funct. Anal. **225** (2008) 2709–2731.
- [CF13] R. Curto, L. Fialkow, *Recursively determined representing measures for bivariate truncated moment sequences*, J. Operator Theory **70(2)** (2013) 401–436.
- [CFM08] R. Curto, L. Fialkow, H. M. Möller, *The extremal truncated moment problem*, Integral Equations Operator Theory **60(2)** (2008) 177–200.
- [CY14] R. Curto, S. Yoo, *Cubic column relations in the truncated moment problems*, J. Funct. Anal. **266(3)** (2014) 1611–1626.
- [CY15] R. Curto, S. Yoo, *Non-extremal sextic moment problems*, J. Funct. Anal. **269(3)** (2015) 758–780.
- [CY16] R. Curto, S. Yoo, *Concrete solution to the nonsingular quartic binary moment problem*, Proc. Amer. Math. Soc. **144** (2016) 249–258.
- [Dan92] J. Dancis, *Positive semidefinite completions of partial hermitian matrices*, Linear Algebra Appl. **175** (1992) 97–114.
- [DS18] P.J. di Dio, K. Schmüdgen. *The multidimensional truncated Moment Problem: Atoms, Determinacy, and Core Variety*, J. Funct. Anal. **274** (2018) 3124–3148. Available from: <https://doi.org/10.1016/j.jfa.2017.11.013>.
- [Fia95] L. Fialkow, *Positivity, extensions and the truncated complex moment problem*, Contemporary Math. **185** (1995), 133–150.
- [Fia11] L. Fialkow, *Solution of the truncated moment problem with variety $y = x^3$* , Trans. Amer. Math. Soc. **363** (2011) 3133–3165.
- [Fia15] L. Fialkow, *The truncated moment problem on parallel lines*, In: Theta Foundation International Book Series of Mathematical Texts **20** (2015), 99–116.
- [Fia17] L. Fialkow. *The core variety of a multisequence in the truncated moment problem*, J. Math. Anal. Appl. **456** (2017) 946–969. Available from: <https://doi.org/10.1016/j.jmaa.2017.07.041>.
- [FN10] L. Fialkow, J. Nie, *Positivity of Riesz functionals and solutions of quadratic and quartic moment problems*, J. Funct. An. **258** (2010), 328–356.
- [GJSW84] R. Grone, C. R. Johnson, E. M. Sá, H. Wolkowicz, *Positive definite completions of partial hermitian matrices*, Linear Algebra Appl. **58** (1984), 109–124.
- [Kim14] D.P. Kimsey. *The cubic complex moment problem*, Integral Equations Operator Theory **80** (2014) 353–378. Available from: <https://doi.org/10.1007/s00020-014-2183-4>.
- [Kim21] D.P. Kimsey. *On a minimal solution for the indefinite truncated multidimensional moment problem*, J. Math. Anal. Appl. **500** (2021) Available from: <https://doi.org/10.1016/j.jmaa.2021.125091>.
- [KN77] K.G. Krein, A.A. Nudelman. *The Markov moment problem and extremal problems*, Translations of Mathematical Monographs, Amer. Math. Soc.; 1977.
- [Lau05] M. Laurent, *Revising two theorems of Curto and Fialkow on moment matrices*, Proc. Amer. Math. Soc. **133** (2005), 2965–2976.
- [Nie14] J. Nie. *The A -truncated K -moment problem*, Found. Comput. Math. **14** (2014) 1243–1276.
- [Sch17] K. Schmüdgen. *The moment problem*, Graduate Texts in Mathematics 277. Cham: Springer; 2017.
- [Wol] Wolfram Research, Inc., Mathematica, Version 12.0, Wolfram Research, Inc., Champaign, IL, 2020.
- [Yoo17a] S. Yoo, *Sextic moment problems on 3 parallel lines*, Bull. Korean Math. Soc. **54** (2017), 299–318.
- [Yoo17b] S. Yoo, *Sextic moment problems with a reducible cubic column relation*, Integral Equations Operator Theory **88** (2017), 45–63.
- [YZ+] S. Yoo, A. Zalar, *The truncated moment problem on reducible cubic curves II: Hyperbolic type relations*, in preparation.
- [Zal21] A. Zalar. *The truncated Hamburger moment problem with gaps in the index set*, Integral Equations Operator Theory **93** (2021) 36 pp. Available from: <https://doi.org/10.1007/s00020-021-02628-6>.
- [Zal22a] A. Zalar. *The truncated moment problem on the union of parallel lines*, Linear Algebra and its Applications **649** (2022) 186–239. Available from: <https://doi.org/10.1016/j.laa.2022.05.008>.

- [Zal22b] A. Zalar. *The strong truncated Hamburger moment problem with and without gaps*, J. Math. Anal. Appl. **516** (2022) 21pp. Available from: <https://doi.org/10.1016/j.jmaa.2022.126563>.
- [Zal23] A. Zalar. *The truncated moment problem on curves $y = q(x)$ and $yx^\ell = 1$* , Linear and Multilinear Algebra (2023) 45pp. Available from: <https://doi.org/10.1080/03081087.2023.2212316>.
- [Zha05] F. Zhang, *The Schur Complement and Its Applications*, Springer-Verlag, New York, 2005.

DEPARTMENT OF MATHEMATICS EDUCATION AND RINS, GYEONGSANG NATIONAL UNIVERSITY, JINJU, 52828, KOREA.

Email address: seyoo@gnu.ac.kr

ALJAŽ ZALAR, FACULTY OF COMPUTER AND INFORMATION SCIENCE, UNIVERSITY OF LJUBLJANA & FACULTY OF MATHEMATICS AND PHYSICS, UNIVERSITY OF LJUBLJANA & INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS, LJUBLJANA, SLOVENIA.

Email address: aljaz.zalar@fri.uni-lj.si